

Summary Report

2018 Community Health Needs Assessment Report

LaSalle Parish

Prepared for:
The Rapides Foundation

By:
Professional Research Consultants, Inc.
11326 P Street Omaha, NE 68137-2316
www.PRCCustomResearch.com

2017-0618-02
© October 2018



Professional Research Consultants, Inc.

Table of Contents

Introduction	4
About This Assessment	5
Methodology	5
PRC Community Health Survey	5
Online Key Informant Survey	7
Public Health, Vital Statistics & Other Data	8
Benchmark Data	9
Determining Significance	10
Information Gaps	10
Summary of Findings	11
Key Data Findings	12
Perceptions of Key Informants	15
Significant Trends	16
Summary Tables	17
Comparisons With Benchmark Data	17
Summary of Key Informant Perceptions	30
Data Charts & Key Informant Input	31
Community Characteristics	32
Population Characteristics	32
Social Determinants of Health	34
General Health Status	38
Overall Health Status	38
Mental Health	42
Death, Disease, & Chronic Conditions	49
Leading Causes of Death	49
Cardiovascular Disease	50
Cancer	57
Respiratory Disease	65
Injury & Violence	68
Diabetes	73
Alzheimer's Disease	77
Kidney Disease	79
Potentially Disabling Conditions	81

Infectious Disease	84
Influenza & Pneumonia Vaccination	85
HIV	87
Sexually Transmitted Diseases	89
Births	91
Birth Outcomes & Risks	91
Family Planning	94
Modifiable Health Risks	96
Nutrition, Physical Activity, & Weight	96
Substance Abuse	107
Tobacco Use	113
Access to Health Services	119
Lack of Health Insurance Coverage (Age 18 to 64)	119
Difficulties Accessing Healthcare	121
Primary Care Services	124
Emergency Room Utilization	126
Oral Health	128
Vision Care	130
Local Resources	131
Perceptions of Local Healthcare Services	131
Quality of Life	132
Life in Central Louisiana	132
Charitable Contribution	133
Civic Participation	134

Introduction



Professional Research Consultants, Inc.

About This Assessment

This Community Health Needs Assessment, a follow-up to similar studies conducted in 2002, 2005, 2010, and 2013, is a systematic, data-driven approach to determining the health status, behaviors and needs of residents in LaSalle Parish, as part of a larger study conducted by The Rapides Foundation. Subsequently, this information may be used to inform decisions and guide efforts to improve community health and wellness.

A Community Health Needs Assessment provides information so that communities may identify issues of greatest concern and decide to commit resources to those areas, thereby making the greatest possible impact on community health status.

This assessment was conducted on behalf of The Rapides Foundation by Professional Research Consultants, Inc. (PRC). PRC is a nationally-recognized healthcare consulting firm with extensive experience conducting Community Health Needs Assessments in hundreds of communities across the United States since 1994.

Methodology

This assessment incorporates data from both quantitative and qualitative sources. Quantitative data input includes primary research (the PRC Community Health Survey) and secondary research (vital statistics and other existing health-related data); these quantitative components allow for comparison to benchmark data at the state and national levels. Qualitative data input includes primary research gathered through an Online Key Informant Survey of various community stakeholders.


PRC Community Health Survey

Survey Instrument

The survey instrument used for this study is based largely on the Centers for Disease Control and Prevention (CDC) Behavioral Risk Factor Surveillance System (BRFSS), as well as various other public health surveys and customized questions addressing gaps in indicator data relative to health promotion and disease prevention objectives and other recognized health issues. The final survey instrument was developed by The Rapides Foundation and PRC.

Community Defined for This Assessment

The focus of the data presented in this report is LaSalle Parish, Louisiana.



Sample Approach & Design

A precise and carefully executed methodology is critical in asserting the validity of the results gathered in the PRC Community Health Survey. Thus, to ensure the best representation of the population surveyed, a mixed-mode methodology was implemented. This included surveys conducted via telephone (landline and cell phone), as well as through online questionnaires.

The sample design used for this effort included a random sample of 196 individuals age 18 and older in LaSalle Parish. All administration of the surveys, data collection and data analysis were conducted by PRC.


For statistical purposes, the maximum rate of error associated with a sample size of 196 respondents is $\pm 7.0\%$ at the 95 percent confidence level.

Sample Characteristics

To accurately represent the population studied, PRC strives to minimize bias through application of a proven telephone methodology and random-selection techniques. While this random sampling of the population produces a highly representative sample, it is a common and preferred practice to “weight” the raw data to improve this representativeness even further. This is accomplished by adjusting the results of a random sample to match the geographic distribution and demographic characteristics of the population surveyed (poststratification), so as to eliminate any naturally occurring bias.

The following chart outlines the characteristics of the LaSalle Parish sample for key demographic variables, compared to actual population characteristics revealed in census data. [Note that the sample consisted solely of area residents age 18 and older; data on children were given by proxy by the person most responsible for that child’s healthcare needs, and these children are not represented demographically in this chart.]

Population & Survey Sample Characteristics (LaSalle Parish, 2018)



Sources: • 2011-2015 American Community Survey, US Census Bureau.
• PRC Community Health Survey, Professional Research Consultants, Inc.

Further note that the poverty descriptions and segmentation used in this report are based on administrative poverty thresholds determined by the US Department of Health & Human Services. These guidelines define poverty status by household income level and number of persons in the household (*e.g., the 2018 guidelines place the poverty threshold for a family of four at \$25,100 annual household income or lower*). In sample segmentation: **“low income”** refers to community members living in a household with defined poverty status or living just above the poverty level and earning up to twice (<200% of) the poverty threshold; **“mid/high income”** refers to those households living on incomes which are twice or more ($\geq 200\%$ of) the federal poverty level.

The sample design and the quality control procedures used in the data collection ensure that the sample is representative. Thus, the findings may be generalized to the total population of community members in the defined area with a high degree of confidence.

Online Key Informant Survey

To solicit input from key informants, those individuals who have a broad interest in the health of the community, an Online Key Informant Survey also was implemented as part of this process. A list of recommended participants was provided by The Rapides Foundation; this list included names and contact information for physicians, public health representatives, other health professionals, social service providers, and a variety of other community leaders. Potential participants were chosen because of their ability to identify primary concerns of the populations with whom they work, as well as of the community overall.

Key informants were contacted by email, introducing the purpose of the survey and providing a link to take the survey online; reminder emails were sent as needed to increase participation. In all, 20 community stakeholders in LaSalle Parish (and 1 who works more broadly throughout Central Louisiana) took part in the Online Key Informant Survey. Final participation included representatives of the organizations in the following list:

- [Ace Family Hardware](#)
- [Hardtner Medical Center](#)
- [Ironhead Youth Development Program](#)
- [LaSalle Association for the Developmentally Delayed](#)
- [LaSalle Economic Development District](#)
- [LaSalle General Hospital](#)
- [LaSalle Parish Library](#)
- [LaSalle Parish Recreation District No. 10](#)
- [LaSalle Parish School Board](#)
- [Northwest Louisiana Human Services District](#)
- [Town of Jena](#)
- [Town of Olla](#)
- [Town of Tullos](#)

Through this process, input was gathered from several individuals whose organizations work with low-income, minority, or other medically underserved populations.

In the online survey, key informants were asked to rate the degree to which various health issues are a problem in their own community. Follow-up questions asked them to describe why they identify problem areas as such, and how these might better be addressed. Results of their ratings, as well as their verbatim comments, are included throughout this report as they relate to the various other data presented.

NOTE: These findings represent qualitative rather than quantitative data. The Online Key Informant Survey was designed to gather input regarding participants' opinions and perceptions of the health needs of the residents in the area. Thus, these findings are not necessarily based on fact.

Public Health, Vital Statistics & Other Data

A variety of existing (secondary) data sources was consulted to complement the research quality of this Community Health Needs Assessment. Data for LaSalle Parish were obtained from the following sources (specific citations are included with the graphs throughout this report):

- [Center for Applied Research and Environmental Systems \(CARES\)](#)
- [Centers for Disease Control & Prevention, Office of Infectious Disease, National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention](#)
- [Centers for Disease Control & Prevention, Office of Public Health Science Services, Center for Surveillance, Epidemiology and Laboratory Services, Division of Health Informatics and Surveillance \(DHIS\)](#)
- [Centers for Disease Control & Prevention, Office of Public Health Science Services, National Center for Health Statistics](#)
- [Community Commons](#)
- [ESRI ArcGIS Map Gallery](#)
- [Louisiana Department of Health](#)

- National Cancer Institute, State Cancer Profiles
- OpenStreetMap (OSM)
- US Census Bureau, American Community Survey
- US Census Bureau, County Business Patterns
- US Census Bureau, Decennial Census
- US Department of Agriculture, Economic Research Service
- US Department of Health & Human Services
- US Department of Health & Human Services, Health Resources and Services Administration (HRSA)
- US Department of Justice, Federal Bureau of Investigation
- US Department of Labor, Bureau of Labor Statistics

Benchmark Data

Trending

A similar survey was administered in LaSalle Parish in 2002, 2005, 2010, and 2013 by PRC on behalf of The Rapides Foundation. Trending data, as revealed by comparison to prior survey results, are provided throughout this report whenever available. Historical data for secondary data indicators are also included for the purposes of trending.

Regional Data

Because this LaSalle Parish survey was part of a larger project covering much of Central Louisiana, comparisons can also be made for many indicators to the broader Rapides Foundation Service Area (referred to as the “RFSA” throughout this report). The RFSA is composed of data from nine Louisiana parishes: Allen, Avoyelles, Catahoula, Grant, LaSalle, Natchitoches, Rapides, Vernon, and Winn.

Louisiana Risk Factor Data

Statewide risk factor data are provided where available as an additional benchmark against which to compare local survey findings; these data are reported in the most recent *BRFSS (Behavioral Risk Factor Surveillance System) Prevalence and Trend Data* published by the Centers for Disease Control and Prevention and the US Department of Health & Human Services. State-level vital statistics are also provided for comparison of secondary data indicators.

Nationwide Risk Factor Data

Nationwide risk factor data, which are also provided in comparison charts, are taken from the *2017 PRC National Health Survey*; the methodological approach for the national study is similar to that employed in this assessment, and these data may be generalized to the US population with a high degree of confidence. National-level vital statistics are also provided for comparison of secondary data indicators.

Healthy People 2020

Healthy People provides science-based, 10-year national objectives for improving the health of all Americans. The Healthy People initiative is grounded in the principle that setting national objectives and monitoring progress can motivate action. For three decades, Healthy People has established benchmarks and monitored progress over time in order to:

- Encourage collaborations across sectors.
- Guide individuals toward making informed health decisions.
- Measure the impact of prevention activities.



Healthy People 2020 is the product of an extensive stakeholder feedback process that is unparalleled in government and health. It integrates input from public health and prevention experts, a wide range of federal, state and local government officials, a consortium of more than 2,000 organizations, and perhaps most importantly, the public. More than 8,000 comments were considered in drafting a comprehensive set of Healthy People 2020 objectives.

Determining Significance

Differences noted in this report represent those determined to be significant. For survey-derived indicators (which are subject to sampling error), statistical significance is determined based on confidence intervals (at the 95 percent confidence level) using question-specific samples and response rates. For the purpose of this report, "significance," of secondary data indicators (which do not carry sampling error but might be subject to reporting error) is determined by a 5% variation from the comparative measure.

Information Gaps

While this assessment is quite comprehensive, it cannot measure all possible aspects of health in the community, nor can it adequately represent all possible populations of interest. It must be recognized that these information gaps might in some ways limit the ability to assess all of the community's health needs.

For example, certain population groups — such as the homeless, institutionalized persons, or those who only speak a language other than English or Spanish — are not represented in the survey data. Other population groups — for example, pregnant women, lesbian/gay/bisexual/transgender residents, undocumented residents, and members of certain racial/ethnic or immigrant groups — might not be identifiable or might not be represented in numbers sufficient for independent analyses.

In terms of content, this assessment was designed to provide a comprehensive and broad picture of the health of the overall community. However, there are certainly medical conditions that are not specifically addressed.

Summary of Findings



Professional Research Consultants, Inc.

Key Data Findings

This summary presents key findings from the data collected for LaSalle Parish for the 2018 Community Health Needs Assessment sponsored by The Rapides Foundation. These include data collected through a (phone and internet-based) random sample population survey, an internet-based survey of key informants, and a review of existing public health data.

Highlighted are differences found when comparing to national data, as well as changes that have occurred since a similar survey was first implemented in LaSalle Parish in 2002.

HEALTH STATUS



Self-Reported Health Status. 20.5% of LaSalle Parish adults characterize their overall health as “fair” or “poor” — this represents a significant decrease from the 31.9% first measured in LaSalle Parish in 2002.

Activity Limitations. 26.9% of LaSalle Parish adults are limited in some way in some activities because of a health-related issue.

Mental Health. 38.6% of LaSalle Parish residents have experienced bouts of depression lasting two or more years during their lives, while 19.1% have been diagnosed with a depressive disorder by a healthcare professional. Overall, 35.3% of the population have ever sought help for mental health, and 4.2% report a time in the past year when they needed such services but were unable to get them.

DEATH & CHRONIC DISEASE



Causes of Death. Cardiovascular disease (heart disease and stroke) and cancers are leading causes of death in LaSalle Parish. Compared to US rates, age-adjusted death rates for many leading causes of death are generally higher in LaSalle Parish (including lung disease, pneumonia/influenza, unintentional injuries [motor vehicle accidents, firearms], diabetes, Alzheimer’s disease, kidney disease, and suicide).

Cancer. A total of 9.6% of adults have ever been diagnosed with cancer.

Heart Disease & Stroke. 10.3% of LaSalle Parish adults report having heart disease (a significant decrease from 16.6% in 2002). Another 6.3% have ever suffered from a stroke.

Diabetes. A total of 18.4% of LaSalle Parish adults have been diagnosed with diabetes.

Lung Disease. 17.6% of LaSalle Parish residents have been diagnosed with chronic obstructive pulmonary disease (which includes chronic bronchitis and emphysema), a prevalence that is significantly above what is found nationally (8.6%).

Overweight & Obesity. Based on reported heights and weights, three-quarters (75.4%) of LaSalle Parish adults are overweight, including 37.3% who are obese. The prevalence of obesity in the parish remains similar to that found in 2002 (30.4%).

INFANT HEALTH & FAMILY PLANNING



Birth Outcomes. Of all births in LaSalle Parish, 8.1% are low-weight (under 2,500g). Additionally, the parish experiences an infant mortality rate of 6.3 deaths for every 1,000 live births (deaths of infants before their first birthday).

Teen Births. The teen birth rate in LaSalle Parish is high, with 57.2 births to girls age 15-19 for every 1,000 girls in this age group (nationally, the teen birth rate is 36.6).

INJURY & VIOLENCE



Unintentional Injury. Death rates due to unintentional injuries (including motor vehicle-related deaths) are much higher than reported nationally (a rate of 81.4 versus 43.7 nationally).

Violence. Rates of violent crime are considerably better in LaSalle Parish than they are nationwide; still, 3.2% of LaSalle Parish adults report experiencing violent crime in the area in the past five years, and 14.6% report ever experiencing domestic violence.

MODIFIABLE HEALTH RISKS



Nutrition. Only four in 10 LaSalle Parish adults (40.2%) get the recommended 5 or more servings of fruits and vegetables per day; however, this is similar to what is found nationally (33.5%). It is important to note that 15.0% of parish adults report difficulty getting fresh produce, and 30.3% say they “sometimes” or “often” ran out of food in the past year before having money to buy more (versus 18.0% nationally).

Physical Activity. Currently, only 17.2% of LaSalle Parish adults meet physical activity guidelines (comparable to the 22.8% nationally). Further, 27.5% of parish adults report not engaging in any type of physical activity outside of work in the month before the survey interview.

Blood Pressure & Cholesterol. In comparison to the nation, LaSalle Parish exhibits a significantly high proportion of adults reporting high blood pressure (51.9% versus 37.0% across the US). A total of 35.1% of parish adults report having high blood cholesterol.

Tobacco Use. 14.9% of LaSalle Parish adults currently smoke cigarettes, much lower than found seen in 2002 (24.6%). Another 9.9% use smokeless tobacco, a prevalence that is higher than the 4.4% nationally and the 3.9% in 2002. A total of 4.4% use electronic cigarettes or vaping devices.

Cardiovascular Risk. A high percentage of LaSalle Parish adults (87.1%) present one or more risk factors or behaviors for heart disease and stroke (including smoking, not getting physical activity, being overweight, or having high blood pressure or cholesterol).

Substance Use. Regarding alcohol use, 12.7% of parish adults are considered to be “binge drinkers,” having had a high number of drinks on a single occasion during the past month (significantly better than the nation’s 20.0%). None of the adults in the parish adults reported illicit drug use in the past month (use of illegal drugs or improper use of prescription drugs), which is significantly below the national prevalence of 2.5%. A total of 22.4% have used prescription opiates (either legally or illegally) in the past year.

PREVENTION



Routine Medical Care. Most parish adults (73.1%) have been to a doctor or clinic for a routine checkup in the past year.

Cancer Screenings. Cancer screening levels in LaSalle Parish are fairly good, including for: *female breast cancer* (75.0% of women age 50-74 have had a mammogram in the past 2 years, compared to 77.0% nationally) and *cervical cancer* (81.0% of women age 21-65 have had a Pap smear in the past 3 years, compared to 73.5% nationally). However, the local proportion of all adults age 50-75 with appropriate colorectal cancer screenings is significantly lower than national levels (66.2% versus 76.4%).

Dental Care. 65.0% of adults in LaSalle Parish have received dental care in the past year. This is much higher than reported in LaSalle Parish in 2002 (47.6%).

Vision Care. A total of 58.1% of LaSalle Parish adults have had a comprehensive eye exam in the past two years, also an improvement since 2002 (48.0%).

ACCESS



Health Insurance Coverage. A total of 10.3% of LaSalle Parish adults between the ages of 18 and 64 are without any type of insurance coverage for health care, either through private or public sources. This is significant improvement from what was recorded in 2002 (19.7%). Still, cost remains a barrier preventing residents from getting medical care (16.7% said they did not get needed medical care in the past year because of the cost).

Difficulties/Delays in Accessing Health Care. A total of 40.8% of LaSalle Parish adults have experienced some type of difficulty or delay in receiving health care in the past year. Difficulty finding physicians and cost are the barriers impacting the greatest shares of adults in LaSalle Parish.

Cost of Prescriptions. A total of 18.8% of LaSalle Parish adults have gone without a needed prescription in the past year because they could not afford it.

Emergency Room Utilization. The proportion of LaSalle Parish adults who have used a local emergency room more than once in the past year (14.5%).

Perceptions of Key Informants

In an online survey of key informants in the area (e.g., a public health professional, physicians, other health providers, social services representatives, community leaders), the following health issues were most often characterized as “major problems” for LaSalle Parish:

- Substance Abuse (86.4% said this is a “major problem” in LaSalle Parish)
- Mental Health (81.8% “major problem”)
- Tobacco Use (72.7% “major problem”)
- Diabetes (68.2% “major problem”)
- Heart Disease and Stroke (54.5% “major problem”)

Significant Trends

The following tables highlight both positive and negative trends observed among the health indicators assessed in this project in comparison with baseline data.

- **Survey Data Indicators:** Trends for survey-derived indicators represent significant changes since 2002 (or 2005, 2010, or 2013 for questions not asked in earlier years).
- **Other Data Indicators:** Trends for other indicators (e.g., public health data) represent point-to-point changes between the most current reporting period and the earliest presented in this report (typically representing the span of 10 to 15 years).

	○ FAVORABLE TRENDS	⬆️ UNFAVORABLE TRENDS
Access to Healthcare Services	<ul style="list-style-type: none"> • Lack of Healthcare Coverage • Specific Source of Ongoing Care 	<ul style="list-style-type: none"> • Difficulty Finding Physician
Cancer	<ul style="list-style-type: none"> • Cancer Deaths 	
Heart Disease & Stroke	<ul style="list-style-type: none"> • Prevalence of Heart Disease • Heart Disease Deaths • Stroke Deaths • Taking Action to Control High Blood Cholesterol • Cardiovascular Risk Factors 	
Injury & Violence	<ul style="list-style-type: none"> • Use of Seat Belts (Adults) 	
Mental Health		<ul style="list-style-type: none"> • “Fair/Poor” Mental Health
Nutrition, Overweight & Physical Activity	<ul style="list-style-type: none"> • Fruit & Vegetable Consumption • Overweights Trying to Lose Weight (Adults) • Medical Advice on Weight • “Fair/Poor” Local Physical Activity Opportunities 	<ul style="list-style-type: none"> • Healthy Weight (Adults) • Overweight (Adults)
Oral Health	<ul style="list-style-type: none"> • Dental Visits (Adults) 	
Potentially Disabling Conditions	<ul style="list-style-type: none"> • “Fair/Poor” Overall Health • [18+] Arthritis/Rheumatism • Eye Exams 	
Respiratory Diseases		<ul style="list-style-type: none"> • CLRD Deaths
Substance Abuse		<ul style="list-style-type: none"> • Drinking & Driving • Rode with Drunk Driver
Tobacco Use	<ul style="list-style-type: none"> • Cigarette Smoking 	<ul style="list-style-type: none"> • Use Smokeless Tobacco

Summary Tables

Comparisons With Benchmark Data

The following tables provide an overview of indicators in LaSalle Parish. These data are grouped to correspond with the Focus Areas presented in Healthy People 2020.






Reading the Data Summary Tables
















■ In the following tables, LaSalle Parish results are shown in the larger, blue column. *Tip: Indicator labels beginning with a “%” symbol are taken from the PRC Community Health Survey; the remaining indicators are taken from secondary data sources.*















■ The columns to the right of the LaSalle Parish column provide trending comparisons (trending from the earliest data year available), as well as comparisons between local data and any available regional (RFSA), state (LA), and national findings, as well as Healthy People 2020 targets. Symbols indicate whether LaSalle Parish compares favorably (☀️), unfavorably (☔️), or comparably (☁️) to these external data.

Note that blank table cells signify that data are not available or are not reliable for that area and/or for that indicator.

Social Determinants	LaSalle Parish	LaSalle Parish vs. Benchmarks				TREND
		vs. RFSA	vs. ST	vs. US	vs. HP2020	
Linguistically Isolated Population (Percent)	0.0	☀️ 0.8	☀️ 1.6	☀️ 4.5		
Population in Poverty (Percent)	23.1	☁️ 20.7	☁️ 19.7	☔️ 15.1		
Population Below 200% FPL (Percent)	45.0	☁️ 44.2	☁️ 39.8	☔️ 33.6		
Children Below 200% FPL (Percent)	46.9	☁️ 52.9	☁️ 49.9	☁️ 43.3		
No High School Diploma (Age 25+, Percent)	24.2	☔️ 18.0	☔️ 16.2	☔️ 13.0		
Unemployment Rate (Age 16+, Percent)	4.2	☀️ 4.9	☁️ 4.3	☁️ 4.1		
% Displaced From Housing in Past 2 Years	7.5	☀️ 13.5			☁️ 7.9	
% "Fair/Poor" Availability of Affordable Housing	47.0	☁️ 45.7			☁️ 40.7	

Social Determinants (continued)	LaSalle Parish	LaSalle Parish vs. Benchmarks				TREND
		vs. RFSA	vs. ST	vs. US	vs. HP2020	
% "Fair/Poor" Condition of Neighborhood Homes	18.7	 20.3				 14.5
						
		better	similar	worse		


























Overall Health	LaSalle Parish	LaSalle Parish vs. Benchmarks				TREND
		vs. RFSA	vs. ST	vs. US	vs. HP2020	
% "Fair/Poor" Overall Health	20.5	 23.3	 21.9	 18.1		 31.9
% 3+ Days Poor Physical Health in Past Month	25.6	 35.4				
% Activity Limitations	26.9	 29.6	 23.0	 25.0		 30.0
% [Limited Activities] Impairment Is Work-Related	24.0	 21.8				
% 4+ Days Health Prevented Usual Activities	18.5	 22.0				 17.7
						
		better	similar	worse		







Access to Health Services	LaSalle Parish	LaSalle Parish vs. Benchmarks				TREND
		vs. RFSA	vs. ST	vs. US	vs. HP2020	
% [Age 18-64] Lack Health Insurance	10.3	 9.2	 16.7	 13.7	 0.0	 19.7
% Difficulty Accessing Healthcare in Past Year (Composite)	40.8	 40.1		 43.2		 34.5
% Difficulty Finding Physician in Past Year	18.8	 14.9		 13.4		 7.7
% Difficulty Getting Appointment in Past Year	13.2	 16.0		 17.5		 11.0











Access to Health Services (continued)	LaSalle Parish	LaSalle Parish vs. Benchmarks				TREND
		vs. RFSA	vs. ST	vs. US	vs. HP2020	
% Cost Prevented Physician Visit in Past Year	16.7	16.6	17.6	15.4		12.9
% Transportation Hindered Dr Visit in Past Year	10.0	9.7		8.3		10.0
% Inconvenient Hrs Prevented Dr Visit in Past Year	8.7	12.4		12.5		9.3
% Cost Prevented Getting Prescription in Past Year	18.8	16.6		14.9		18.8
Primary Care Doctors per 100,000	47.2	58.7	78.7	87.8		
% Have a Specific Source of Ongoing Care	79.7	76.7		74.1	95.0	67.8
% Have Had Routine Checkup in Past Year	73.1	77.0	72.1	68.3		78.1
% Two or More ER Visits in Past Year	14.5	12.9		9.3		15.0
% Rate Local Healthcare "Fair/Poor"	14.1	19.1		16.2		






better similar worse
















































Cancer	LaSalle Parish	LaSalle Parish vs. Benchmarks				TREND
		vs. RFSA	vs. ST	vs. US	vs. HP2020	
Cancer (Age-Adjusted Death Rate)	153.4	184.4	179.4	158.5	161.4	191.2
Female Breast Cancer Incidence Rate	80.4	106.3	123.2	123.5		
Prostate Cancer Incidence Rate	127.4	140.9	144.4	114.8		
Lung Cancer Incidence Rate	82.5	76.0	70.5	61.2		

Cancer (continued)	LaSalle Parish	LaSalle Parish vs. Benchmarks				TREND
		vs. RFSA	vs. ST	vs. US	vs. HP2020	
Colorectal Cancer Incidence Rate	39.1	 52.3	 47.8	 39.8		
% Cancer	9.6	 8.5				 6.3
% [Women 50-74] Mammogram in Past 2 Years	75.0	 79.8	 78.5	 77.0	 81.1	 68.9
% [Women 21-65] Pap Smear in Past 3 Years	81.0	 76.1	 81.5	 73.5	 93.0	 74.7
% [Men 50+] Prostate Exam in Past 2 Years	74.7	 70.1				 78.3
% [Age 50-75] Colorectal Cancer Screening	66.2	 74.9	 64.1	 76.4	 70.5	 67.7
						
		better	similar	worse		

Dementias, Including Alzheimer's Disease	LaSalle Parish	LaSalle Parish vs. Benchmarks				TREND
		vs. RFSA	vs. ST	vs. US	vs. HP2020	
Alzheimer's Disease (Age-Adjusted Death Rate)	39.8	 48.7	 37.7	 26.6		
						
		better	similar	worse		

Diabetes	LaSalle Parish	LaSalle Parish vs. Benchmarks				TREND
		vs. RFSA	vs. ST	vs. US	vs. HP2020	
Diabetes (Age-Adjusted Death Rate)	29.4	 24.1	 27.1	 21.4	 20.5	
% Diabetes/High Blood Sugar	18.4	 16.3	 12.1	 13.3		 13.9
% Borderline/Pre-Diabetes	5.3	 7.3		 9.5		

Diabetes (continued)	LaSalle Parish	LaSalle Parish vs. Benchmarks				TREND
		vs. RFSA	vs. ST	vs. US	vs. HP2020	
% [Non-Diabetes] Blood Sugar Tested in Past 3 Years	62.1	 53.9		 50.0		
						
		better	similar	worse		

Heart Disease & Stroke	LaSalle Parish	LaSalle Parish vs. Benchmarks				TREND
		vs. RFSA	vs. ST	vs. US	vs. HP2020	
Diseases of the Heart (Age-Adjusted Death Rate)	142.9	 244.7	 213.8	 167.0	 156.9	 321.1
Stroke (Age-Adjusted Death Rate)	40.1	 55.9	 45.9	 37.1	 34.8	 75.8
% Heart Disease (Heart Attack, Angina, Coronary Disease)	10.3	 8.8		 8.0		 16.6
% Stroke	6.3	 4.5	 4.0	 4.7		 9.2
% Blood Pressure Checked in Past 2 Years	97.8	 94.2		 90.4	 92.6	 98.5
% Told Have High Blood Pressure (Ever)	51.9	 46.7	 39.3	 37.0	 26.9	 50.8
% [HBP] Taking Action to Control High Blood Pressure	92.2	 92.4		 93.8		 87.0
% Cholesterol Checked in Past 5 Years	84.8	 87.2	 77.7	 85.1	 82.1	 85.7
% Told Have High Cholesterol (Ever)	35.1	 35.1		 36.2	 13.5	 34.1
% [HBC] Taking Action to Control High Blood Cholesterol	95.8	 90.5		 87.3		 74.4
% 1+ Cardiovascular Risk Factor	87.1	 91.3		 87.2		 96.4
						
		better	similar	worse		



























HIV	LaSalle Parish	LaSalle Parish vs. Benchmarks				TREND
		vs. RFSA	vs. ST	vs. US	vs. HP2020	
HIV Prevalence Rate	662.6	369.6	502.3	353.2	22.1	
		better	similar	worse		








Immunization & Infectious Diseases	LaSalle Parish	LaSalle Parish vs. Benchmarks				TREND
		vs. RFSA	vs. ST	vs. US	vs. HP2020	
% [Age 65+] Flu Vaccine in Past Year	72.0	70.9	51.6	76.8	70.0	74.6
% [Age 65+] Pneumonia Vaccine Ever	79.5	71.8	73.1	82.7	90.0	73.1
		better	similar	worse		

Infant Health & Family Planning	LaSalle Parish	LaSalle Parish vs. Benchmarks				TREND
		vs. RFSA	vs. ST	vs. US	vs. HP2020	
Low Birthweight Births (Percent)	8.1	10.3	10.9	8.2	7.8	8.3
Infant Death Rate	6.3	7.6	8.4	6.2	6.0	
Teen Births per 1,000 (Age 15-19)	57.2	60.9	50.2	36.6		65.3
		better	similar	worse		

Injury & Violence	LaSalle Parish	LaSalle Parish vs. Benchmarks				TREND
		vs. RFSA	vs. ST	vs. US	vs. HP2020	
Unintentional Injury (Age-Adjusted Death Rate)	81.4	57.3	54.0	43.7	36.4	72.8
Motor Vehicle Crashes (Age-Adjusted Death Rate)	36.2	21.5	17.5	10.9	12.4	
% "Always" Wear Seat Belt	83.1	83.1				67.2
Firearm-Related Deaths (Age-Adjusted Death Rate)	13.8	15.2	19.3	10.5	9.3	
Violent Crime Rate	311.2	590.3	512.9	379.7		
% Victim of Violent Crime in Past 5 Years	3.2	3.3		3.7		0.8
% Victim of Domestic Violence (Ever)	14.6	17.4		14.2		11.3
% Victim of Domestic Violence in Past 5 Years	8.0	5.6				
		better	similar	worse		

Kidney Disease	LaSalle Parish	LaSalle Parish vs. Benchmarks				TREND
		vs. RFSA	vs. ST	vs. US	vs. HP2020	
Kidney Disease (Age-Adjusted Death Rate)	25.3	22.9	25.2	13.9		
		better	similar	worse		

Mental Health	LaSalle Parish	LaSalle Parish vs. Benchmarks				TREND
		vs. RFSA	vs. ST	vs. US	vs. HP2020	
% "Fair/Poor" Mental Health	19.1	 18.6		 13.0		 9.9
% 3+ Days Poor Mental Health in Past Month	31.0	 33.0				
% Diagnosed Depression	19.1	 27.9	 19.9	 21.6		
% Symptoms of Chronic Depression (2+ Years)	38.6	 38.2		 31.4		 40.5
Suicide (Age-Adjusted Death Rate)	14.8	 14.0	 12.8	 12.4	 10.2	
% Had Someone to Turn to "All/Most" of the Time in Past Month	86.7	 81.8				
% Taking Rx/Receiving Mental Health Trtmt	15.8	 21.1		 13.9		 11.7
% Have Ever Sought Help for Mental Health	35.3	 34.1		 30.8		 32.3
% Unable to Get Mental Health Svcs in Past Yr	4.2	 7.3		 6.8		
						
		better	similar	worse		

Nutrition, Physical Activity & Weight	LaSalle Parish	LaSalle Parish vs. Benchmarks				TREND
		vs. RFSA	vs. ST	vs. US	vs. HP2020	
% Eat 5+ Servings of Fruit or Vegetables per Day	40.2	 34.1		 33.5		 25.3
% [Adults] Eats 2+ Servings of Fruit per Day	51.5	 47.9				 38.0
% [Adults] Eats 3+ Servings of Vegetables per Day	34.5	 28.0				 25.1

Nutrition, Physical Activity & Weight (continued)	LaSalle Parish	LaSalle Parish vs. Benchmarks				TREND
		vs. RFSA	vs. ST	vs. US	vs. HP2020	
% Difficulty Getting Fresh Fruits & Vegetables	15.0	15.0				14.4
% Medical Advice About Nutrition in Past Year	39.4	44.0				43.4
% "Often/Sometimes" Ran Out of Food in the Past Year	30.3	32.9		18.0		
Population With Low Food Access (Percent)	28.3	31.5	26.8	22.4		
% No Leisure-Time Physical Activity	27.5	30.3	29.1	26.2	32.6	28.5
% Meeting Physical Activity Guidelines	17.2	19.9	18.7	22.8	20.1	
% [Adults] Vigorous Physical Activity	29.2	29.6				
% [Adults] Moderate Physical Activity	21.9	23.7				
% Strengthening Activity	25.5	28.6	27.2	33.8		19.1
% Walk Regularly (5+ Times Per Week For >10 Minutes)	39.6	40.1				42.3
% "Often" See Others in Community Being Physically Active	42.7	39.7				39.5
% "Fair/Poor" Local Physical Activity Opportunities	33.1	36.5				41.3
Recreation/Fitness Facilities per 100,000	6.7	4.8	9.5	11.0		
% Medical Advice About Exercise in Past Year	35.9	44.4				41.3
% Overweight (BMI 25+)	75.4	75.3	69.2	67.8		66.4
% Healthy Weight (BMI 18.5-24.9)	23.4	23.4	29.0	30.3	33.9	32.1

Nutrition, Physical Activity & Weight (continued)	LaSalle Parish	LaSalle Parish vs. Benchmarks				TREND
		vs. RFSA	vs. ST	vs. US	vs. HP2020	
% [Overweights] Trying to Lose Weight Both Diet/Exercise	41.1	34.7				20.7
% Obese (BMI 30+)	37.3	41.4	35.5	32.8	30.5	30.4
% Medical Advice on Weight in Past Year	29.0	26.6		24.2		17.3
% [Overweights] Counseled About Weight in Past Year	32.4	32.3		29.0		
		better	similar	worse		












Oral Health	LaSalle Parish	LaSalle Parish vs. Benchmarks				TREND
		vs. RFSA	vs. ST	vs. US	vs. HP2020	
% [Age 18+] Dental Visit in Past Year	65.0	53.6	56.6	59.7	49.0	47.6
		better	similar	worse		


























Potentially Disabling Conditions	LaSalle Parish	LaSalle Parish vs. Benchmarks				TREND
		vs. RFSA	vs. ST	vs. US	vs. HP2020	
% [50+] Arthritis/Rheumatism	48.7	45.6		38.3		
% [18+] Arthritis/Rheumatism	29.5	28.7		23.1		46.5
% Eye Exam in Past 2 Years	58.1	58.2		55.3		48.0
		better	similar	worse		











Respiratory Diseases	LaSalle Parish	LaSalle Parish vs. Benchmarks				TREND
		vs. RFSA	vs. ST	vs. US	vs. HP2020	
CLRD (Age-Adjusted Death Rate)	74.8	56.3	43.9	40.9		57.5
Pneumonia/Influenza (Age-Adjusted Death Rate)	40.2	26.5	16.6	14.8		
% COPD (Lung Disease)	17.6	14.7	8.3	8.6		15.4
		better	similar	worse		

Sexually Transmitted Diseases	LaSalle Parish	LaSalle Parish vs. Benchmarks				TREND
		vs. RFSA	vs. ST	vs. US	vs. HP2020	
Chlamydia Incidence Rate	351.9	536.0	625.9	456.1		
Gonorrhea Incidence Rate	81.2	154.6	194.6	110.7		
		better	similar	worse		

Substance Abuse	LaSalle Parish	LaSalle Parish vs. Benchmarks				TREND
		vs. RFSA	vs. ST	vs. US	vs. HP2020	
% Current Drinker	38.1	49.0	51.9	55.0		
% Binge Drinker (Single Occasion - 5+ Drinks Men, 4+ Women)	12.7	22.1	16.9	20.0	24.4	8.1
% Excessive Drinker	13.5	23.6		22.5	25.4	
% Drinking & Driving in Past Month	4.7	3.8	3.5	5.2		1.1
% Rode w/Drunk Driver in Past Month	5.8	5.9				1.1

Substance Abuse (continued)	LaSalle Parish	LaSalle Parish vs. Benchmarks				TREND
		vs. RFSA	vs. ST	vs. US	vs. HP2020	
% Illicit Drug Use in Past Month	0.0	 3.2		 2.5	 7.1	 0.2
% Have Used Prescription Opiates in Past Year	22.4	 25.0				
% Ever Sought Help for Alcohol or Drug Problem	2.4	 4.3		 3.4		 1.9
						
		better	similar	worse		


Tobacco Use	LaSalle Parish	LaSalle Parish vs. Benchmarks				TREND
		vs. RFSA	vs. ST	vs. US	vs. HP2020	
% Current Smoker	14.9	 23.6	 22.8	 16.3	 12.0	 24.6
% Someone Smokes at Home	15.5	 16.6		 10.7		 20.5
% [Nonsmokers] Someone Smokes in the Home	7.8	 7.0		 4.0		
% Aware of Smoking Cessation Services/Programs	25.7	 41.2				 28.6
% Community Believes Adults "Definitely" Should Not Smoke	43.1	 41.2				 49.3
% Currently Use Vaping Products	4.4	 5.6	 6.0	 3.8		
% Use Smokeless Tobacco	9.9	 7.2	 5.1	 4.4	 0.3	 3.9
						
		better	similar	worse		

Quality of Life	LaSalle Parish	LaSalle Parish vs. Benchmarks				TREND
		vs. RFSA	vs. ST	vs. US	vs. HP2020	
% "Fair/Poor" Overall Quality of Life in Central Louisiana	19.4	 28.2				
% Parish Life: Wrong Track and Getting Worse	7.1	 17.0				
% Know 10+ People Benefiting from Charities	44.3	 40.0				
% "Frequently/Sometimes" Donate to Charity	67.6	 67.4				
% Have Received Charitable Assistance in Past Year	8.5	 6.1				
% "Frequently/Sometimes" Volunteer	36.1	 40.9				
% Voted in Each of the Past 5 Elections	65.4	 54.5				
		 better	 similar	 worse		

Summary of Key Informant Perceptions

In the Online Key Informant Survey, community stakeholders were asked to rate the degree to which each of 20 health issues is a problem in their own community, using a scale of “major problem,” “moderate problem,” “minor problem,” or “no problem at all.” The following chart summarizes their responses; these findings also are outlined throughout this report, along with the qualitative input describing reasons for their concerns. (Note that these ratings alone do not establish priorities for this assessment; rather, they are one of several data inputs considered for the prioritization process described earlier.)

Key Informants: Relative Position of Health Topics as Problems in the Community



Data Charts & Key Informant Input

The following sections present data from multiple sources, including the random-sample PRC Community Health Survey, public health and other existing data sets (secondary data), as well as qualitative input from the Online Key Informant Survey. Data indicators from these sources are intermingled and organized by health topic. To better understand the source data for specific indicators, please refer to the footnotes accompanying each chart.



Professional Research Consultants, Inc.

Community Characteristics

Population Characteristics

Land Area, Population Size & Density

Data from the US Census Bureau reveal the following statistics for our community relative to size, population, and density.

Total Population
(Estimated Population, 2012-2016)

	Total Population	Total Land Area (Square Miles)	Population Density (Per Square Mile)
LaSalle Parish	14,927	624.68	23.9
Louisiana	4,645,670	43,206.73	107.52
United States	318,558,162	3,532,068.58	90.19


Sources: • US Census Bureau American Community Survey 5-year estimates.
• Retrieved August 2018 from Community Commons at <http://www.chna.org>.

Age

It is important to understand the age distribution of the population, as different age groups have unique health needs that should be considered separately from others along the age spectrum.

- LaSalle Parish has a higher proportion of seniors compared to Louisiana and the US.

Total Population by Age Groups, Percent
(2012-2016)




Sources: • US Census Bureau American Community Survey 5-year estimates.
• Retrieved August 2018 from Community Commons at <http://www.chna.org>.

Race & Ethnicity

The following charts illustrate the racial and ethnic makeup of our community. Note that ethnicity (Hispanic or Latino) can be of any race.

- LaSalle Parish is racially less diverse than the state or nation.


Total Population by Race Alone, Percent
(2012-2016)



Sources: • US Census Bureau American Community Survey 5-year estimates.
• Retrieved August 2018 from Community Commons at <http://www.chna.org>.

- The LaSalle Parish Hispanic proportion is similar to the state but well below the US.

Hispanic Population
(2012-2016)



Sources: • US Census Bureau American Community Survey 5-year estimates.
• Retrieved August 2018 from Community Commons at <http://www.chna.org>.
Notes: • Origin can be viewed as the heritage, nationality group, lineage, or country of birth of the person or the person's parents or ancestors before their arrival in the United States. People who identify their origin as Hispanic, Latino, or Spanish may be of any race.

Social Determinants of Health

About Social Determinants


Health starts in our homes, schools, workplaces, neighborhoods, and communities. We know that taking care of ourselves by eating well and staying active, not smoking, getting the recommended immunizations and screening tests, and seeing a doctor when we are sick all influence our health. Our health is also determined in part by access to social and economic opportunities; the resources and supports available in our homes, neighborhoods, and communities; the quality of our schooling; the safety of our workplaces; the cleanliness of our water, food, and air; and the nature of our social interactions and relationships. The conditions in which we live explain in part why some Americans are healthier than others and why Americans more generally are not as healthy as they could be.

- Healthy People 2020 (www.healthypeople.gov)

Poverty

The following chart outlines the proportion of our population below the federal poverty threshold, as well as below 200% of the federal poverty level, in comparison to state and national proportions.

- The proportion of the parish population living in poverty (below 100% and also 200% of the federal level) is similar to the state but higher than the nation.




- Sources:
- US Census Bureau American Community Survey 5-year estimates.
 - Retrieved August 2018 from Community Commons at <http://www.chna.org>.
- Notes:
- Poverty is considered a key driver of health status. This indicator is relevant because poverty creates barriers to access including health services, healthy food, and other necessities that contribute to poor health status.

Education

Education levels are reflected in the proportion of our population without a high school diploma:

- The proportion of LaSalle Parish adults without a high school education is higher than found in the Rapides Foundation Service Area, Louisiana, and the US.

Population With No High School Diploma (Population Age 25+ Without a High School Diploma or Equivalent, 2012-2016)



Sources: • US Census Bureau American Community Survey 5-year estimates.
 • Retrieved August 2018 from Community Commons at <http://www.chna.org>.
 Notes: • This indicator is relevant because educational attainment is linked to positive health outcomes.

Housing


Survey respondents were asked:

“Overall, how would you rate the availability of affordable housing in your community? Would you say: excellent, very good, good, fair, or poor?”

“How would you describe the condition of the homes in your neighborhood? Would you say: excellent, very good, good, fair, or poor?”


- Adults in LaSalle Parish and the Rapides Foundation Service Area rate affordable housing availability and the condition of local housing similarly.
- Though fluctuating considerably, “air/poor” responses regarding affordability and the condition of housing in LaSalle Parish have not changed significantly compared to 2002 survey findings.

Perceive the Availability of Affordable Local Housing to be “Fair” or “Poor”



Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 336]
 Notes: • Asked of all respondents.

Perceive the Condition of Neighborhood Homes to be “Fair” or “Poor”




Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 338]
 Notes: • Asked of all respondents.

“Because of an emergency, have you had to live with a friend or relative in the past two years, even if this was only temporary?”

- The current parish finding is more favorable than found throughout the Rapides Foundation Service Area and is comparable to the 2002 baseline.

Had to Live With a Friend/Relatives in the Past Two Years Due to an Emergency (Even if Only Temporarily)



Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 337]
 Notes: • Asked of all respondents.

General Health Status


Overall Health Status

Self-Reported Health Status

The initial inquiry of the PRC Community Health Survey asked respondents the following:

“Would you say that in general your health is: excellent, very good, good, fair, or poor?”

Self-Reported Health Status (LaSalle Parish, 2018)




Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 5]
Notes: • Asked of all respondents.

The following charts further detail “fair/poor” overall health responses in LaSalle Parish in comparison to past findings and benchmark data, as well as by basic demographic characteristics (namely by sex, age groupings, and income [based on poverty status]).

- “Fair/poor” evaluations of overall health in LaSalle Parish are similar to what is found regionally (RFSA), statewide, and nationally; these evaluations have improved considerably since the baseline 2002 assessment.


Experience “Fair” or “Poor” Overall Health



Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 5]
 • Behavioral Risk Factor Surveillance System Survey Data. Atlanta, Georgia. United States Department of Health and Human Services, Centers for Disease Control and Prevention (CDC): 2016 LA data.
 • 2017 PRC National Health Survey, Professional Research Consultants, Inc.
 Notes: • Asked of all respondents.

- “Fair/poor” health evaluations in LaSalle Parish are considerably higher among women and adults age 65 and older. The difference by income level is not statistically significant.

Experience “Fair” or “Poor” Overall Health (LaSalle Parish, 2018)



Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 5]
 Notes: • Asked of all respondents.
 • Income categories reflect respondent’s household income as a ratio to the federal poverty level (FPL) for their household size. “Low Income” includes households with incomes up to 200% of the federal poverty level; “Mid/High Income” includes households with incomes at 200% or more of the federal poverty level.

Activity Limitations

About Disability & Health

An individual can get a disabling impairment or chronic condition at any point in life. Compared with people without disabilities, people with disabilities are more likely to:

- Experience difficulties or delays in getting the health care they need.
- Not have had an annual dental visit.
- Not have had a mammogram in past 2 years.
- Not have had a Pap test within the past 3 years.
- Not engage in fitness activities.
- Use tobacco.
- Be overweight or obese.
- Have high blood pressure.
- Experience symptoms of psychological distress.
- Receive less social-emotional support.
- Have lower employment rates.


There are many social and physical factors that influence the health of people with disabilities. The following three areas for public health action have been identified, using the International Classification of Functioning, Disability, and Health (ICF) and the three World Health Organization (WHO) principles of action for addressing health determinants.

- **Improve the conditions of daily life** by: encouraging communities to be accessible so all can live in, move through, and interact with their environment; encouraging community living; and removing barriers in the environment using both physical universal design concepts and operational policy shifts.
- **Address the inequitable distribution of resources among people with disabilities and those without disabilities** by increasing: appropriate health care for people with disabilities; education and work opportunities; social participation; and access to needed technologies and assistive supports.
- **Expand the knowledge base and raise awareness about determinants of health for people with disabilities** by increasing: the inclusion of people with disabilities in public health data collection efforts across the lifespan; the inclusion of people with disabilities in health promotion activities; and the expansion of disability and health training opportunities for public health and health care professionals.
- Healthy People 2020 (www.healthypeople.gov)

“Are you limited in any way in any activities because of physical, mental, or emotional problems?”

- The proportion of parish adults with activity limitations is similar to the region, state, and nation. The parish rate is statistically unchanged since the 2002 baseline study.


Limited in Activities in Some Way Due to a Physical, Mental or Emotional Problem



Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 128]
 • Behavioral Risk Factor Surveillance System Survey Data. Atlanta, Georgia. United States Department of Health and Human Services, Centers for Disease Control and Prevention (CDC): 2015 LA data.
 • 2017 PRC National Health Survey, Professional Research Consultants, Inc.
 Notes: • Asked of all respondents.

- Activity limitations are considerably higher among women than men. Differences by age and income level are not statistically significant.

Limited in Activities in Some Way Due to a Physical, Mental or Emotional Problem (LaSalle Parish, 2018)



Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 109]
 Notes: • Asked of all respondents.
 • Income categories reflect respondent's household income as a ratio to the federal poverty level (FPL) for their household size. "Low Income" includes households with incomes up to 200% of the federal poverty level; "Mid/High Income" includes households with incomes at 200% or more of the federal poverty level.

Mental Health

About Mental Health & Mental Disorders

Mental health is a state of successful performance of mental function, resulting in productive activities, fulfilling relationships with other people, and the ability to adapt to change and to cope with challenges. Mental health is essential to personal well-being, family and interpersonal relationships, and the ability to contribute to community or society. Mental disorders are health conditions that are characterized by alterations in thinking, mood, and/or behavior that are associated with distress and/or impaired functioning. Mental disorders contribute to a host of problems that may include disability, pain, or death. Mental illness is the term that refers collectively to all diagnosable mental disorders. Mental disorders are among the most common causes of disability. The resulting disease burden of mental illness is among the highest of all diseases.

Mental health and physical health are closely connected. Mental health plays a major role in people's ability to maintain good physical health. Mental illnesses, such as depression and anxiety, affect people's ability to participate in health-promoting behaviors. In turn, problems with physical health, such as chronic diseases, can have a serious impact on mental health and decrease a person's ability to participate in treatment and recovery.

The existing model for understanding mental health and mental disorders emphasizes the interaction of social, environmental, and genetic factors throughout the lifespan. In behavioral health, researchers identify: **risk factors**, which predispose individuals to mental illness; and **protective factors**, which protect them from developing mental disorders. Researchers now know that the prevention of mental, emotional, and behavioral (MEB) disorders is inherently interdisciplinary and draws on a variety of different strategies. Over the past 20 years, research on the prevention of mental disorders has progressed. The major areas of progress include evidence that:


- MEB disorders are common and begin early in life.
- The greatest opportunity for prevention is among young people.
- There are multiyear effects of multiple preventive interventions on reducing substance abuse, conduct disorder, antisocial behavior, aggression, and child maltreatment.
- The incidence of depression among pregnant women and adolescents can be reduced.
- School-based violence prevention can reduce the base rate of aggressive problems in an average school by 25 to 33%.
- There are potential indicated preventive interventions for schizophrenia.
- Improving family functioning and positive parenting can have positive outcomes on mental health and can reduce poverty-related risk.
- School-based preventive interventions aimed at improving social and emotional outcomes can also improve academic outcomes.
- Interventions targeting families dealing with adversities, such as parental depression or divorce, can be effective in reducing risk for depression in children and increasing effective parenting.
- Some preventive interventions have benefits that exceed costs, with the available evidence strongest for early childhood interventions.
- Implementation is complex, and it is important that interventions be relevant to the target audiences.
- In addition to advancements in the prevention of mental disorders, there continues to be steady progress in treating mental disorders as new drugs and stronger evidence-based outcomes become available.

- Healthy People 2020 (www.healthypeople.gov)

Self-Reported Mental Health Status

“Now thinking about your mental health, which includes stress, depression, and problems with emotions, would you say that, in general, your mental health is: excellent, very good, good, fair, or poor?”


Self-Reported Mental Health Status (LaSalle Parish, 2018)



Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 99]
 Notes: • Asked of all respondents.

- The “fair/poor” response in LaSalle Parish is higher than the national finding.
- Evaluations of “fair” or “poor” mental health are higher among adults ages 18 to 64 and those at lower incomes. There is no significant difference by gender.

Experience “Fair” or “Poor” Mental Health (LaSalle Parish, 2018)




Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 99]
 Notes: • Asked of all respondents.
 • Income categories reflect respondent’s household income as a ratio to the federal poverty level (FPL) for their household size. “Low Income” includes households with incomes up to 200% of the federal poverty level; “Mid/High Income” includes households with incomes at 200% or more of the federal poverty level.

Depression

Diagnosed Depression: “Has a doctor or other healthcare provider ever told you that you have a depressive disorder, including depression, major depression, dysthymia, or minor depression?”

- The proportion of parish adults with diagnosed depression is lower than found in the service area but is similar to the state and nation.

Have Been Diagnosed With a Depressive Disorder




- Sources:
- 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 119]
 - Behavioral Risk Factor Surveillance System Survey Data. Atlanta, Georgia. United States Department of Health and Human Services, Centers for Disease Control and Prevention (CDC): 2016 LA data.
 - 2017 PRC National Health Survey, Professional Research Consultants, Inc.
- Notes:
- Asked of all respondents.
 - Depressive disorders include depression, major depression, dysthymia, or minor depression.

Symptoms of Chronic Depression: “Have you had two years or more in your life when you felt depressed or sad most days, even if you felt okay sometimes?”

- Symptoms of chronic depression in the parish are statistically similar to the service area and US. The parish rate is comparable to the 2002 baseline, though fluctuating considerably over time.


Have Experienced Symptoms of Chronic Depression



- Sources:
- 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 117]
 - 2017 PRC National Health Survey, Professional Research Consultants, Inc.
- Notes:
- Asked of all respondents.
 - Chronic depression includes periods of two or more years during which the respondent felt depressed or sad on most days, even if (s)he felt okay sometimes.

- Prevalence of chronic depression is much higher among adults age 18 to 64 and lower-income adults. There is no significant difference by gender.

Have Experienced Symptoms of Chronic Depression (LaSalle Parish, 2018)




Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 100]
 Notes: • Asked of all respondents.
 • Chronic depression includes periods of two or more years during which the respondent felt depressed or sad on most days, even if (s)he felt okay sometimes.
 • Income categories reflect respondent's household income as a ratio to the federal poverty level (FPL) for their household size. "Low Income" includes households with incomes up to 200% of the federal poverty level; "Mid/High Income" includes households with incomes at 200% or more of the federal poverty level.

Emotional Support

“In the past month, how often have you had someone you could turn to if you needed or wanted help? Would you say: all of the time, most of the time, some of the time, little of the time, or none of the time?”

- Levels of emotional support are comparable across the parish and Rapides Foundation Service Area.

Had Someone to Turn to “All” or “Most” of the Time in the Past Month (LaSalle Parish, 2018)




Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 330]
 Notes: • Asked of all respondents.

Suicide

The following chart outlines the most current age-adjusted mortality rates attributed to suicide in our population. (Refer to “Leading Causes of Death” for an explanation of the use of age-adjusting for these rates.)

- The parish suicide rate is statistically similar to the service area and state but worse than the US.

Suicide: Age-Adjusted Mortality Trends
 (2007-2016 Annual Average Deaths per 100,000 Population)
 Healthy People 2020 Target = 10.2 or Lower



Sources: • CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics. Data extracted August 2018.
 • US Department of Health and Human Services. Healthy People 2020. December 2010. <http://www.healthypeople.gov> [Objective MHMD-1]

Notes: • Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10).
 • Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population.


Mental Health Treatment

“Have you ever sought help from a professional for a mental or emotional problem?”

“Are you now taking medication or receiving treatment from a doctor or other health professional for any type of mental health condition or emotional problem?”

- The proportions of parish adults seeking help and taking medication/receiving treatment are comparable to the national findings.

Mental Health Treatment




Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Items 103-104]
 • 2017 PRC National Health Survey, Professional Research Consultants, Inc.
 Notes: • Reflects the total sample of respondents.

“Was there a time in the past 12 months when you needed mental health services but were not able to get them?”

- Adults in the parish and across the nation had similar difficulty obtaining mental health services.
- Within LaSalle Parish, a significant difference was found between income groups.

Unable to Get Mental Health Services When Needed in the Past Year (LaSalle Parish, 2018)



Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 105]
 Notes: • Asked of all respondents.
 • Income categories reflect respondent's household income as a ratio to the federal poverty level (FPL) for their household size. "Low Income" includes households with incomes up to 200% of the federal poverty level; "Mid/High Income" includes households with incomes at 200% or more of the federal poverty level.

Key Informant Input: Mental Health

The following chart outlines key informants' perceptions of the severity of *Mental Health* as a problem in the community:

Perceptions of Mental Health as a Problem in the Community (Key Informants, 2018)

■ Major Problem ■ Moderate Problem ■ Minor Problem ■ No Problem At All



Sources: ● PRC Online Key Informant Survey, Professional Research Consultants, Inc.
Notes: ● Asked of all respondents.

Challenges

Among those rating this issue as a “major problem,” the following represent what key informants see as the main challenges for persons with mental illness:

Access to Care/Services

We are in a rural parish. Both hospitals have behavioral health units for senior adults only. There are no licensed mental health clinics available for adults. There are some services being provided to youth in the schools, outside contracted companies. – Community Leader (LaSalle Parish)

No nearby mental health clinics or specialists in the area. – Social Services Provider (LaSalle Parish)


Death, Disease, & Chronic Conditions

Leading Causes of Death

Distribution of Deaths by Cause

Cancers and cardiovascular disease (heart disease and stroke) are leading causes of death in the community.

Leading Causes of Death
(LaSalle Parish, 2014-2016)



- Sources: • CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics. Data extracted August 2018.
- Notes: • Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10).
• CLRD is chronic lower respiratory disease.

Age-Adjusted Death Rates for Selected Causes

In order to compare mortality in the region with other localities (in this case, the state and the United States), it is necessary to look at *rates* of death — these are figures which represent the number of deaths in relation to the population size (such as deaths per 100,000 population, as is used here).

Furthermore, in order to compare localities without undue bias toward younger or older populations, the common convention is to adjust the data to some common baseline age distribution. Use of these “age-adjusted” rates provides the most valuable means of gauging mortality against benchmark data, as well as *Healthy People 2020* targets.

Charts throughout this report outline annual average age-adjusted death rates per 100,000 population for selected causes of death in the area. (For infant mortality data, see also *Birth Outcomes & Risks* in the **Births** section of this report.)

Cardiovascular Disease

About Heart Disease & Stroke

Heart disease is the leading cause of death in the United States, with stroke following as the third leading cause. Together, heart disease and stroke are among the most widespread and costly health problems facing the nation today, accounting for more than \$500 billion in healthcare expenditures and related expenses in 2010 alone. Fortunately, they are also among the most preventable.

The leading modifiable (controllable) risk factors for heart disease and stroke are:

- High blood pressure
- High cholesterol
- Cigarette smoking
- Diabetes
- Poor diet and physical inactivity
- Overweight and obesity

The risk of Americans developing and dying from cardiovascular disease would be substantially reduced if major improvements were made across the US population in diet and physical activity, control of high blood pressure and cholesterol, smoking cessation, and appropriate aspirin use.

The burden of cardiovascular disease is disproportionately distributed across the population. There are significant disparities in the following based on gender, age, race/ethnicity, geographic area, and socioeconomic status:

- Prevalence of risk factors
- Access to treatment
- Appropriate and timely treatment
- Treatment outcomes
- Mortality

Disease does not occur in isolation, and cardiovascular disease is no exception. Cardiovascular health is significantly influenced by the physical, social, and political environment, including: maternal and child health; access to educational opportunities; availability of healthy foods, physical education, and extracurricular activities in schools; opportunities for physical activity, including access to safe and walkable communities; access to healthy foods; quality of working conditions and worksite health; availability of community support and resources; and access to affordable, quality healthcare.

- Healthy People 2020 (www.healthypeople.gov)

Age-Adjusted Heart Disease & Stroke Deaths


The greatest share of cardiovascular deaths is attributed to heart disease. The following charts outline age-adjusted mortality rates for heart disease and for stroke in our community.

- Heart disease deaths in LaSalle Parish have decreased significantly over time and have trended much lower than state and national rates.

Heart Disease: Age-Adjusted Mortality Trends

(Annual Average Deaths per 100,000 Population)

Healthy People 2020 Target = 156.9 or Lower (Adjusted)



Sources: • CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics. Data extracted August 2018.
 • US Department of Health and Human Services. Healthy People 2020. December 2010. <http://www.healthypeople.gov> [Objective HDS-2]


Notes: • Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10).
 • Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population.
 • The Healthy People 2020 Heart Disease target is adjusted to account for all diseases of the heart.

- Stroke deaths in the parish have declined significantly to comparable levels with the state and US.

Stroke: Age-Adjusted Mortality Trends

(Annual Average Deaths per 100,000 Population)

Healthy People 2020 Target = 34.8 or Lower



Sources: • CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics. Data extracted August 2018.
 • US Department of Health and Human Services. Healthy People 2020. December 2010. <http://www.healthypeople.gov> [Objective HDS-3]

Notes: • Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10).
 • Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population.


Prevalence of Heart Disease & Stroke

“Has a doctor, nurse, or other health professional ever told you that you had: a heart attack, also called a myocardial infarction; or angina or coronary heart disease?” (Heart disease prevalence here is a calculated prevalence that includes those responding affirmatively to either.)

“Has a doctor, nurse, or other health professional ever told you that you had a stroke?”

- Prevalence of heart disease in the parish is similar to the region and nation. Over time, prevalence of heart disease has improved within the parish.


Prevalence of Heart Disease



Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 146]
 • 2017 PRC National Health Survey, Professional Research Consultants, Inc.
 Notes: • Asked of all respondents.
 • Includes diagnoses of heart attack, angina or coronary heart disease.

- Prevalence of stroke in the parish is similar to all populations shown in the following chart. The parish rate remains statistically unchanged over time.

Prevalence of Stroke



Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 35]
 • 2017 PRC National Health Survey, Professional Research Consultants, Inc.
 • Behavioral Risk Factor Surveillance System Survey Data. Atlanta, Georgia. United States Department of Health and Human Services, Centers for Disease Control and Prevention (CDC); 2016 LA data.
 Notes: • Asked of all respondents.

Cardiovascular Risk Factors

About Cardiovascular Risk

Controlling risk factors for heart disease and stroke remains a challenge. High blood pressure and cholesterol are still major contributors to the national epidemic of cardiovascular disease. High blood pressure affects approximately 1 in 3 adults in the United States, and more than half of Americans with high blood pressure do not have it under control. High sodium intake is a known risk factor for high blood pressure and heart disease, yet about 90% of American adults exceed their recommendation for sodium intake.

- Healthy People 2020 (www.healthypeople.gov)

High Blood Pressure & Cholesterol Prevalence

“Have you ever been told by a doctor, nurse, or other health care professional that you had high blood pressure?”


“Are you currently taking any action to help control your high blood pressure, such as taking medication, changing your diet, or exercising?”

“Blood cholesterol is a fatty substance found in the blood. Have you ever been told by a doctor, nurse, or other health care professional that your blood cholesterol is high?”

“Are you currently taking any action to help control your high cholesterol, such as taking medication, changing your diet, or exercising?”

- Prevalence of high blood pressure in LaSalle Parish is comparable to the service area but is less favorable than found across the state and nation. After a decline, prevalence in the parish has returned to the baseline level.


Prevalence of High Blood Pressure Healthy People 2020 Target = 26.9% or Lower



Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Items 41, 129]
 • Behavioral Risk Factor Surveillance System Survey Data. Atlanta, Georgia. United States Department of Health and Human Services, Centers for Disease Control and Prevention (CDC): 2015 LA data.
 • 2017 PRC National Health Survey, Professional Research Consultants, Inc.
 • US Department of Health and Human Services. Healthy People 2020. December 2010. <http://www.healthypeople.gov> [Objective HDS-5.1]
 Notes: • Asked of all respondents.

- High cholesterol prevalence in LaSalle Parish is identical to the service area and similar to the US. Parish trending has not varied statistically since 2002.

Prevalence of High Blood Cholesterol Healthy People 2020 Target = 13.5% or Lower



Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Items 46, 148]
 • 2017 PRC National Health Survey, Professional Research Consultants, Inc.
 • US Department of Health and Human Services. Healthy People 2020. December 2010. <http://www.healthypeople.gov> [Objective HDS-7]
 Notes: • Asked of all respondents.

About Cardiovascular Risk

Individual level risk factors which put people at increased risk for cardiovascular diseases include:

- High Blood Pressure
- High Blood Cholesterol
- Tobacco Use
- Physical Inactivity
- Poor Nutrition
- Overweight/Obesity
- Diabetes
- National Center for Chronic Disease Prevention and Health Promotion, Centers for Disease Control and Prevention

Three health-related behaviors contribute markedly to cardiovascular disease:

Poor nutrition. People who are overweight have a higher risk for cardiovascular disease. Almost 60% of adults are overweight or obese. To maintain a proper body weight, experts recommend a well-balanced diet which is low in fat and high in fiber, accompanied by regular exercise.

Lack of physical activity. People who are not physically active have twice the risk for heart disease of those who are active. More than half of adults do not achieve recommended levels of physical activity.

Tobacco use. Smokers have twice the risk for heart attack of nonsmokers. Nearly one-fifth of all deaths from cardiovascular disease, or about 190,000 deaths a year nationally, are smoking-related. Every day, more than 3,000 young people become daily smokers in the US.

Modifying these behaviors is critical both for preventing and for controlling cardiovascular disease. Other steps that adults who have cardiovascular disease should take to reduce their risk of death and disability include adhering to treatment for high blood pressure and cholesterol, using aspirin as appropriate, and learning the symptoms of heart attack and stroke.


- National Center for Chronic Disease Prevention and Health Promotion, Centers for Disease Control and Prevention

Total Cardiovascular Risk

The following chart reflects the percentage of adults in the LaSalle Parish who report one or more of the following: being overweight; smoking cigarettes; being physically inactive; or having high blood pressure or cholesterol. See also *Nutrition, Physical Activity, Weight Status, and Tobacco Use* in the **Modifiable Health Risks** section of this report.

- Cardiovascular risk is nearly identical across the parish and nation.
- Within the parish, seniors and lower-income adults are much more likely to have cardiovascular risk factors.

Present One or More Cardiovascular Risks or Behaviors
(LaSalle Parish, 2018)




Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 131]
 Notes: • Asked of all respondents.
 • Cardiovascular risk is defined as exhibiting one or more of the following: 1) no leisure-time physical activity; 2) regular/occasional cigarette smoking; 3) hypertension; 4) high blood cholesterol; and/or 5) being overweight/obese.
 • Income categories reflect respondent's household income as a ratio to the federal poverty level (FPL) for their household size. "Low Income" includes households with incomes up to 200% of the federal poverty level; "Mid/High Income" includes households with incomes at 200% or more of the federal poverty level.

Key Informant Input: Heart Disease & Stroke

The following chart outlines key informants' perceptions of the severity of *Heart Disease & Stroke* as a problem in the community:

Perceptions of Heart Disease and Stroke as a Problem in the Community
(Key Informants, 2018)



Sources: • PRC Online Key Informant Survey, Professional Research Consultants, Inc.
 Notes: • Asked of all respondents.

Top Concerns

Among those rating this issue as a “major problem,” reasons related to the following:

Prevalence/Incidence

So many people in our community suffer from heart disease. There seems to be very little knowledge about how to avoid it, and obesity runs rampant. – Community Leader (LaSalle Parish)

They affected almost every family in our community. I feel diet and exercise play major roles in this. – Community Leader (LaSalle Parish)

Access to Care/Services

Same barriers as with cancer and diabetes. Not enough physicians with expertise to care for the disease. Not enough patient education to help them properly manage the disease. Barriers to physical activity and proper nutrition education. – Community Leader (LaSalle Parish)

Cancer

About Cancer

Continued advances in cancer research, detection, and treatment have resulted in a decline in both incidence and death rates for all cancers. Among people who develop cancer, more than half will be alive in five years. Yet, cancer remains a leading cause of death in the United States, second only to heart disease.

Many cancers are preventable by reducing risk factors such as: use of tobacco products; physical inactivity and poor nutrition; obesity; and ultraviolet light exposure. Other cancers can be prevented by getting vaccinated against human papillomavirus and hepatitis B virus. In the past decade, overweight and obesity have emerged as new risk factors for developing certain cancers, including colorectal, breast, uterine corpus (endometrial), and kidney cancers. The impact of the current weight trends on cancer incidence will not be fully known for several decades. Continued focus on preventing weight gain will lead to lower rates of cancer and many chronic diseases.

Screening is effective in identifying some types of cancers (see US Preventive Services Task Force [USPSTF] recommendations), including:


- Breast cancer (using mammography)
- Cervical cancer (using Pap tests)
- Colorectal cancer (using fecal occult blood testing, sigmoidoscopy, or colonoscopy)
- Healthy People 2020 (www.healthypeople.gov)

Age-Adjusted Cancer Deaths

The following chart illustrates age-adjusted cancer mortality (all types) in LaSalle Parish.

- Cancer deaths in the parish are lower than the Louisiana rate but similar to the US rate. Over time, the parish death rate has declined significantly.

Cancer: Age-Adjusted Mortality Trends
(Annual Average Deaths per 100,000 Population)
Healthy People 2020 Target = 161.4 or Lower



Sources: • CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics. Data extracted August 2018.
• US Department of Health and Human Services. Healthy People 2020. December 2010. <http://www.healthypeople.gov> [Objective C-1]

Notes: • Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10).
• Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population.

Cancer Risk

About Cancer Risk

Reducing the nation's cancer burden requires reducing the prevalence of behavioral and environmental factors that increase cancer risk.

- All cancers caused by cigarette smoking could be prevented. At least one-third of cancer deaths that occur in the United States are due to cigarette smoking.
- According to the American Cancer Society, about one-third of cancer deaths that occur in the United States each year are due to nutrition and physical activity factors, including obesity.
- National Center for Chronic Disease Prevention and Health Promotion, Centers for Disease Control and Prevention

Cancer Screenings

The American Cancer Society recommends that both men and women get a cancer-related checkup during a regular doctor's checkup. It should include examination for cancers of the thyroid, testicles, ovaries, lymph nodes, oral cavity, and skin, as well as health counseling about tobacco, sun exposure, diet and nutrition, risk factors, sexual practices, and environmental and occupational exposures.

Screening levels in the community were measured in the PRC Community Health Survey relative to: female breast cancer (mammography); cervical cancer (Pap smear testing); and colorectal cancer (sigmoidoscopy and fecal occult blood testing).

Female Breast Cancer Screening

About Screening for Breast Cancer

The US Preventive Services Task Force (USPSTF) recommends screening mammography, with or without clinical breast examination (CBE), every 1-2 years for women age 40 and older.

Rationale: The USPSTF found fair evidence that mammography screening every 12-33 months significantly reduces mortality from breast cancer. Evidence is strongest for women age 50-69, the age group generally included in screening trials. For women age 40-49, the evidence that screening mammography reduces mortality from breast cancer is weaker, and the absolute benefit of mammography is smaller, than it is for older women. Most, but not all, studies indicate a mortality benefit for women undergoing mammography at ages 40-49, but the delay in observed benefit in women younger than 50 makes it difficult to determine the incremental benefit of beginning screening at age 40 rather than at age 50.


The absolute benefit is smaller because the incidence of breast cancer is lower among women in their 40s than it is among older women. The USPSTF concluded that the evidence is also generalizable to women age 70 and older (who face a higher absolute risk for breast cancer) if their life expectancy is not compromised by comorbid disease. The absolute probability of benefits of regular mammography increase along a continuum with age, whereas the likelihood of harms from screening (false-positive results and unnecessary anxiety, biopsies, and cost) diminish from ages 40-70. The balance of benefits and potential harms, therefore, grows more favorable as women age. The precise age at which the potential benefits of mammography justify the possible harms is a subjective choice. The USPSTF did not find sufficient evidence to specify the optimal screening interval for women age 40-49.

- US Preventive Services Task Force, Agency for Healthcare Research and Quality, US Department of Health & Human Services

Note that other organizations (e.g., American Cancer Society, American Academy of Family Physicians, American College of Physicians, National Cancer Institute) may have slightly different screening guidelines.

Breast Cancer Screening: “A mammogram is an x-ray of each breast to look for cancer. How long has it been since you had your last mammogram?” (Calculated here among women age 50 to 74 who indicate screening within the past 2 years.)

- Parish screenings are similar to all populations shown below and statistically unchanged over time.



Sources:

- 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 151]
- Behavioral Risk Factor Surveillance System Survey Data. Atlanta, Georgia. United States Department of Health and Human Services, Centers for Disease Control and Prevention (CDC): 2016 LA data.
- 2017 PRC National Health Survey, Professional Research Consultants, Inc.
- US Department of Health and Human Services. Healthy People 2020. December 2010. <http://www.healthypeople.gov> [Objective C-17]

Notes:

- Reflects female respondents 50-74.

Cervical Cancer Screenings

About Screening for Cervical Cancer

The US Preventive Services Task Force (USPSTF) strongly recommends screening for cervical cancer in women who have been sexually active and have a cervix.

Rationale: The USPSTF found good evidence from multiple observational studies that screening with cervical cytology (Pap smears) reduces incidence of and mortality from cervical cancer. Direct evidence to determine the optimal starting and stopping age and interval for screening is limited. Indirect evidence suggests most of the benefit can be obtained by beginning screening within 3 years of onset of sexual activity or age 21 (whichever comes first) and screening at least every 3 years. The USPSTF concludes that the benefits of screening substantially outweigh potential harms.

The USPSTF recommends against routinely screening women older than age 65 for cervical cancer if they have had adequate recent screening with normal Pap smears and are not otherwise at high risk for cervical cancer.

Rationale: The USPSTF found limited evidence to determine the benefits of continued screening in women older than 65. The yield of screening is low in previously screened women older than 65 due to the declining incidence of high-grade cervical lesions after middle age. There is fair evidence that screening women older than 65 is associated with an increased risk for potential harms, including false-positive results and invasive procedures. The USPSTF concludes that the potential harms of screening are likely to exceed benefits among older women who have had normal results previously and who are not otherwise at high risk for cervical cancer.

The USPSTF recommends against routine Pap smear screening in women who have had a total hysterectomy for benign disease.

Rationale: The USPSTF found fair evidence that the yield of cytologic screening is very low in women after hysterectomy and poor evidence that screening to detect vaginal cancer improves health outcomes. The USPSTF concludes that potential harms of continued screening after hysterectomy are likely to exceed benefits.

- US Preventive Services Task Force, Agency for Healthcare Research and Quality, US Department of Health & Human Services


Note that other organizations (e.g., American Cancer Society, American Academy of Family Physicians, American College of Physicians, National Cancer Institute) may have slightly different screening guidelines.

Cervical Cancer Screening: “A Pap test is a test for cancer of the cervix. How long has it been since you had your last Pap test?” (Calculated here among women age 21 to 65 who indicate screening within the past 3 years.)

- Parish screenings are comparable to all populations shown in the following chart. The increase over time is not statistically significant.

Have Had a Pap Smear in the Past Three Years (Among Women Age 21-65)

Healthy People 2020 Target = 93.0% or Higher



Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 152]
 • Behavioral Risk Factor Surveillance System Survey Data. Atlanta, Georgia. United States Department of Health and Human Services, Centers for Disease Control and Prevention (CDC); 2016 LA data.
 • 2017 PRC National Health Survey, Professional Research Consultants, Inc.
 • US Department of Health and Human Services. Healthy People 2020. December 2010. <http://www.healthypeople.gov> [Objective C-15]

Notes: • Reflects female respondents age 21 to 65.

Prostate Cancer Screenings

About Screening for Prostate Cancer

Prostate cancer is one of the most common types of cancer that affects men. In the United States, the lifetime risk of being diagnosed with prostate cancer is approximately 11%, and the lifetime risk of dying of prostate cancer is 2.5%. Many men with prostate cancer never experience symptoms and, without screening, would never know they have the disease. In autopsy studies of men who died of other causes, more than 20% of men aged 50 to 59 years and more than 33% of men aged 70 to 79 years were found to have prostate cancer. In some men, the cancer is more aggressive and leads to death. The median age of death from prostate cancer is 80 years, and more than two-thirds of all men who die of prostate cancer are older than 75 years.

Screening for prostate cancer begins with a test that measures the amount of PSA protein in the blood. An elevated PSA level may be caused by prostate cancer but can also be caused by other conditions, including an enlarged prostate (benign prostatic hyperplasia) and inflammation of the prostate (prostatitis). Some men without prostate cancer may therefore have positive screening results (i.e., “false-positive” results). Men with a positive PSA test result may undergo a transrectal ultrasound-guided core-needle biopsy of the prostate to diagnose prostate cancer.

PSA-based screening for prostate cancer has both potential benefits and harms. The USPSTF *does not* recommend screening for prostate cancer unless men express a preference for screening after being informed of and understanding the benefits and risks. The decision about whether to be screened for prostate cancer requires that each man incorporate his own values about the potential benefits and harms. The potential harms of screening, diagnostic procedures, and treatment occur soon after screening takes place. Although the potential benefits may occur any time after screening, they generally occur years after treatment, because progression from asymptomatic, screen-detected cancer to symptomatic, metastasized cancer or death (if it occurs at all) may take years or decades to occur.

The USPSTF concludes with moderate certainty that the net benefit of PSA-based screening for prostate cancer in men aged 55 to 69 years is small for some men. How each man weighs specific benefits and harms will determine whether the overall net benefit is small.

The USPSTF concludes with moderate certainty that the potential benefits of PSA-based screening for prostate cancer in men 70 years and older do not outweigh the expected harms.


- US Preventive Services Task Force, Agency for Healthcare Research and Quality, US Department of Health & Human Services

Note that other organizations (e.g., American Cancer Society, American Academy of Family Physicians, American College of Physicians, National Cancer Institute) may have slightly different screening guidelines.

Prostate Cancer Screening: “A prostate-specific antigen test, also called a PSA test, is a blood test used to check men for prostate cancer. How long has it been since you had your last PSA test?”

“A digital rectal exam is when a doctor, nurse or other health professional places a gloved finger in the rectum to feel the size, hardness and shape of the prostate gland. How long has it been since your last digital rectal exam?” (Calculated here among men age 50 and older who indicate either screening within the past 2 years.)

- Screening levels are comparable across the parish and Rapides Foundation Service Area. The parish rate is statistically similar to the 2002 baseline.



Colorectal Cancer Screenings

About Screening for Colorectal Cancer

The USPSTF recommends screening for colorectal cancer using fecal occult blood testing, sigmoidoscopy, or colonoscopy in adults, beginning at age 50 years and continuing until age 75 years.

The evidence is convincing that screening for colorectal cancer with fecal occult blood testing, sigmoidoscopy, or colonoscopy detects early-stage cancer and adenomatous polyps. There is convincing evidence that screening with any of the three recommended tests (fecal occult blood testing, sigmoidoscopy, colonoscopy) reduces colorectal cancer mortality in adults age 50 to 75 years. Follow-up of positive screening test results requires colonoscopy regardless of the screening test used.

- US Preventive Services Task Force, Agency for Healthcare Research and Quality, US Department of Health & Human Services

Note that other organizations (e.g., American Cancer Society, American Academy of Family Physicians, American College of Physicians, National Cancer Institute) may have slightly different screening guidelines.

Colorectal Cancer Screening: “Sigmoidoscopy and colonoscopy are exams in which a tube is inserted in the rectum to view the colon for signs of cancer or other health problems. How long has it been since your last sigmoidoscopy or colonoscopy?” and

“A blood stool test is a test that may use a special kit at home to determine whether the stool contains blood. How long has it been since you had your last blood stool test?”


(Calculated here among both sexes age 50 to 75 who indicated fecal occult blood testing within the past year and/or sigmoidoscopy/colonoscopy [lower endoscopy] within the past 10 years.)

- Screenings in LaSalle Parish are similar to the state but less favorable than the service area and nation. The parish rate has not varied statistically since 2013.

Have Had a Colorectal Cancer Screening

(Among Adults Age 50-75)

Healthy People 2020 Target = 70.5% or Higher



- Sources:
- 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 155]
 - Behavioral Risk Factor Surveillance System Survey Data. Atlanta, Georgia. United States Department of Health and Human Services, Centers for Disease Control and Prevention (CDC); 2016 LA data.
 - 2017 PRC National Health Survey, Professional Research Consultants, Inc.
 - US Department of Health and Human Services. Healthy People 2020. December 2010. <http://www.healthypeople.gov> [Objective C-16]
- Notes:
- Asked of all respondents age 50 through 75.
 - In this case, the term "colorectal screening" refers to adults age 50-75 receiving a FOBT (fecal occult blood test) in the past year and/or a lower endoscopy (sigmoidoscopy/colonoscopy) in the past 10 years.

Key Informant Input: Cancer

The following chart outlines key informants' perceptions of the severity of *Cancer* as a problem in the community:

Perceptions of Cancer as a Problem in the Community (Key Informants, 2018)

■ Major Problem ■ Moderate Problem ■ Minor Problem ■ No Problem At All



Sources: • PRC Online Key Informant Survey, Professional Research Consultants, Inc.
Notes: • Asked of all respondents.

Top Concerns

Among those rating this issue as a “major problem,” reasons related to the following:

Prevalence/Incidence

It seems that every week I hear about someone new with cancer. In our hospital alone, we have had several ladies with breast cancer over the years, plus children/grandchildren of many of our employees have had cancer, as well as children in our parish. – Other Health Provider (LaSalle Parish)

You can name a member of nearly every family that has been affected by cancer in some way. Treatment for this disease is an hour away, at best, which causes hardships on many families in terms of expense and time. – Other Health Provider (LaSalle Parish)

Cancer rates for this area are extremely high. Oil field work conditions, something in the water, tobacco, drug use, manufacturing plant? Not sure the cause! – Community Leader (LaSalle Parish)

It seems as though many people in our community are stricken with cancer. Locally, we have no diagnostic or treatment centers, counseling centers, etc. – Community Leader (LaSalle Parish)

There are always people mentioning that they have some form of cancer in our parish. – Community Leader (LaSalle Parish)

Many people suffering from cancer; younger people being diagnosed with very serious problems, with some dying at an early age. – Community Leader (LaSalle Parish)

Number of community members with a cancer diagnosis, along with poor lifestyle choices. Healthcare. Reduced physical activity for various reasons, poor diet for various reasons. – Community Leader (LaSalle Parish)

There appears to be an overwhelming number of persons with cancer in our community. – Community Leader (LaSalle Parish)

Because it seems as though everyone around here has it. – Community Leader (LaSalle Parish)

So many people in this community get cancer. – Community Leader (LaSalle Parish)

Access to Care/Services

There seems to be a large number of cancer patients, but there is no cancer treatment facilities anywhere nearby. – Social Services Provider (LaSalle Parish)

Respiratory Disease

About Asthma & COPD

Asthma and chronic obstructive pulmonary disease (COPD) are significant public health burdens. Specific methods of detection, intervention, and treatment exist that may reduce this burden and promote health.

Asthma is a chronic inflammatory disorder of the airways characterized by episodes of reversible breathing problems due to airway narrowing and obstruction. These episodes can range in severity from mild to life threatening. Symptoms of asthma include wheezing, coughing, chest tightness, and shortness of breath. Daily preventive treatment can prevent symptoms and attacks and enable individuals who have asthma to lead active lives.

COPD is a preventable and treatable disease characterized by airflow limitation that is not fully reversible. The airflow limitation is usually progressive and associated with an abnormal inflammatory response of the lung to noxious particles or gases (typically from exposure to cigarette smoke). Treatment can lessen symptoms and improve quality of life for those with COPD.

The burden of respiratory diseases affects individuals and their families, schools, workplaces, neighborhoods, cities, and states. Because of the cost to the healthcare system, the burden of respiratory diseases also falls on society; it is paid for with higher health insurance rates, lost productivity, and tax dollars. Annual healthcare expenditures for asthma alone are estimated at \$20.7 billion.

Asthma. The prevalence of asthma has increased since 1980. However, deaths from asthma have decreased since the mid-1990s. The causes of asthma are an active area of research and involve both genetic and environmental factors.

Risk factors for asthma currently being investigated include:

- Having a parent with asthma
- Sensitization to irritants and allergens
- Respiratory infections in childhood
- Overweight

Asthma affects people of every race, sex, and age. However, significant disparities in asthma morbidity and mortality exist, in particular for low-income and minority populations. Populations with higher rates of asthma include: children; women (among adults) and boys (among children); African Americans; Puerto Ricans; people living in the Northeast United States; people living below the Federal poverty level; and employees with certain exposures in the workplace.

While there is not a cure for asthma yet, there are diagnoses and treatment guidelines that are aimed at ensuring that all people with asthma live full and active lives.

- Healthy People 2020 (www.healthypeople.gov)

[NOTE: COPD was changed to chronic lower respiratory disease (CLRD) with the introduction of ICD-10 codes. CLRD is used in vital statistics reporting, but COPD is still widely used and commonly found in surveillance reports.]


Age-Adjusted Respiratory Disease Deaths

Chronic lower respiratory diseases (CLRD) are diseases affecting the lungs; the most deadly of these is chronic obstructive pulmonary disease (COPD), which includes emphysema and chronic bronchitis.

Pneumonia and influenza mortality also is illustrated in the following chart. For prevalence of vaccinations against pneumonia and influenza, see also *Immunization & Infectious Diseases* in the **Infectious Disease** section of this report.

- The CLRD death rate has trended upward in LaSalle Parish and is much higher than found across the state and nation.

CLRD: Age-Adjusted Mortality Trends (Annual Average Deaths per 100,000 Population)




Sources: CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics. Data extracted August 2018.

Notes: Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10). Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population. CLRD is chronic lower respiratory disease.

- The pneumonia/influenza death rate in LaSalle Parish is much worse than found across the region, state, and nation.

Pneumonia/Influenza: Age-Adjusted Mortality Trends (2012-2016 Annual Average Deaths per 100,000 Population)



Sources: CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics. Data extracted August 2018.


Notes: Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10). Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population.

Prevalence of Respiratory Diseases

COPD

“Would you please tell me if you have ever suffered from or been diagnosed with COPD or chronic obstructive pulmonary disease, including bronchitis or emphysema?”

- COPD prevalence in the parish is similar to the service area but worse than Louisiana and the US. The parish level has not varied statistically since 2002.




Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 24]
 • Behavioral Risk Factor Surveillance System Survey Data. Atlanta, Georgia. United States Department of Health and Human Services, Centers for Disease Control and Prevention (CDC): 2016 LA data.
 • 2017 PRC National Health Survey, Professional Research Consultants, Inc.

Notes: • Asked of all respondents.
 • Includes those having ever suffered from or been diagnosed with COPD or chronic obstructive pulmonary disease, including bronchitis or emphysema.
 • In prior data, the term "chronic lung disease" was used, which also included bronchitis or emphysema.

Key Informant Input: Respiratory Disease

This chart outlines perceptions of the severity of *Respiratory Disease* as a problem in the community:

Perceptions of Respiratory Diseases as a Problem in the Community (Key Informants, 2018)



Sources: • PRC Online Key Informant Survey, Professional Research Consultants, Inc.
 Notes: • Asked of all respondents.

Top Concerns

Among those rating this issue as a “major problem,” the following reason was given:

Tobacco Use

Excessive smoking habits are prevalent in the community. – Other Health Provider (LaSalle Parish)

Injury & Violence

About Injury & Violence

Injuries and violence are widespread in society. Both unintentional injuries and those caused by acts of violence are among the top 15 killers for Americans of all ages. Many people accept them as “accidents,” “acts of fate,” or as “part of life.” However, most events resulting in injury, disability, or death are predictable and preventable.

Injuries are the leading cause of death for Americans ages 1 to 44, and a leading cause of disability for all ages, regardless of sex, race/ethnicity, or socioeconomic status. More than 180,000 people die from injuries each year, and approximately 1 in 10 sustains a nonfatal injury serious enough to be treated in a hospital emergency department.

Beyond their immediate health consequences, injuries and violence have a significant impact on the well-being of Americans by contributing to:

- Premature death
- Disability
- Poor mental health
- High medical costs
- Lost productivity

The effects of injuries and violence extend beyond the injured person or victim of violence to family members, friends, coworkers, employers, and communities.

Numerous factors can affect the risk of unintentional injury and violence, including individual behaviors, physical environment, access to health services (ranging from pre-hospital and acute care to rehabilitation), and social environment (from parental monitoring and supervision of youth to peer group associations, neighborhoods, and communities).

Interventions addressing these social and physical factors have the potential to prevent unintentional injuries and violence.

Efforts to prevent unintentional injury may focus on:

- Modifications of the environment
- Improvements in product safety
- Legislation and enforcement
- Education and behavior change
- Technology and engineering


Efforts to prevent violence may focus on:

- Changing social norms about the acceptability of violence
- Improving problem-solving skills (for example, parenting, conflict resolution, coping)
- Changing policies to address the social and economic conditions that often give rise to violence
- Healthy People 2020 (www.healthypeople.gov)

Leading Causes of Accidental Death

Leading causes of accidental death in the area include the following:

Leading Causes of Accidental Death (LaSalle Parish, 2014-2016)



Sources: • CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics. Data extracted August 2018.
Notes: • Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10).


Unintentional Injury

Age-Adjusted Unintentional Injury Deaths

The following chart outlines age-adjusted mortality rates for unintentional injury in the area.

- The parish rate is worse than the state and national rates but is statistically similar to the baseline rate.

Unintentional Injuries: Age-Adjusted Mortality Trends (Annual Average Deaths per 100,000 Population) Healthy People 2020 Target = 36.4 or Lower




Sources: • CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics. Data extracted August 2018.
• US Department of Health and Human Services. Healthy People 2020. December 2010. <http://www.healthypeople.gov> [Objective IVP-11]
Notes: • Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10).
• Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population.

Intentional Injury (Violence)

Violent Crime

Violent crime is composed of four offenses (FBI Index offenses): murder and non-negligent manslaughter; forcible rape; robbery; and aggravated assault. Note that the quality of crime data can vary widely from location to location, depending on the consistency and completeness of reporting among various jurisdictions.

- The violent crime rate in LaSalle Parish is considerably lower than the service area, state, and nation.




- Sources:
- Federal Bureau of Investigation, FBI Uniform Crime Reports.
 - Retrieved August 2018 from Community Commons at <http://www.chna.org>.
- Notes:
- This indicator reports the rate of violent crime offenses reported by the sheriff's office or county police department per 100,000 residents. Violent crime includes homicide, rape, robbery, and aggravated assault. This indicator is relevant because it assesses community safety.
 - Participation by law enforcement agencies in the UCR program is voluntary. Sub-state data do not necessarily represent an exhaustive list of crimes due to gaps in reporting. Also, some institutions of higher education have their own police departments, which handle offenses occurring within campus grounds; these offenses are not included in the violent crime statistics, but can be obtained from the Uniform Crime Reports Universities and Colleges data tables.

Violent Crime Experience: "Have you been the victim of a violent crime in your area in the past 5 years?"

- The proportion of violent crime victims in the parish is comparable to the national rate.
- There are no significant differences within any of the demographic groups.

Victim of a Violent Crime in the Past Five Years (LaSalle Parish, 2018)




Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 46]
 Notes: • Asked of all respondents.
 • Income categories reflect respondent's household income as a ratio to the federal poverty level (FPL) for their household size. "Low Income" includes households with incomes up to 200% of the federal poverty level; "Mid/High Income" includes households with incomes at 200% or more of the federal poverty level.

Intimate Partner Violence: “The next questions are about different types of violence in relationships with an intimate partner. By an intimate partner, I mean any current or former spouse, boyfriend, or girlfriend. Someone you were dating, or romantically or sexually intimate with, would also be considered an intimate partner. Has an intimate partner ever hit, slapped, pushed, kicked, or hurt you in any way?”

- Prevalence of domestic violence in the parish is comparable to the region and nation. In LaSalle Parish, the prevalence is statistically unchanged over time.

Have Ever Been Hit, Slapped, Pushed, Kicked, or Hurt in Any Way by an Intimate Partner




Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 50]
 • 2017 PRC National Health Survey, Professional Research Consultants, Inc.
 Notes: • Asked of all respondents.

Key Informant Input: Injury & Violence

The following chart outlines key informants' perceptions of the severity of *Injury & Violence* as a problem in the community:

Perceptions of Injury and Violence as a Problem in the Community (Key Informants, 2018)

■ Major Problem ■ Moderate Problem ■ Minor Problem ■ No Problem At All



Sources: • PRC Online Key Informant Survey, Professional Research Consultants, Inc.
Notes: • Asked of all respondents.

Diabetes

About Diabetes

Diabetes mellitus occurs when the body cannot produce or respond appropriately to insulin. Insulin is a hormone that the body needs to absorb and use glucose (sugar) as fuel for the body's cells. Without a properly functioning insulin signaling system, blood glucose levels become elevated and other metabolic abnormalities occur, leading to the development of serious, disabling complications. Many forms of diabetes exist; the three common types are Type 1, Type 2, and gestational diabetes. Effective therapy can prevent or delay diabetic complications.

Diabetes mellitus:

- Lowers life expectancy by up to 15 years.
- Increases the risk of heart disease by 2 to 4 times.
- Is the leading cause of kidney failure, lower limb amputations, and adult-onset blindness.

The rate of diabetes mellitus continues to increase both in the United States and throughout the world. Due to the steady rise in the number of persons with diabetes mellitus, and possibly earlier onset of type 2 diabetes mellitus, there is growing concern about the possibility that the increase in the number of persons with diabetes mellitus and the complexity of their care might overwhelm existing healthcare systems.

People from minority populations are more frequently affected by type 2 diabetes. Minority groups constitute 25% of all adult patients with diabetes in the US and represent the majority of children and adolescents with type 2 diabetes.

Lifestyle change has been proven effective in preventing or delaying the onset of type 2 diabetes in high-risk individuals.

- Healthy People 2020 (www.healthypeople.gov)

Age-Adjusted Diabetes Deaths

Age-adjusted diabetes mortality for the area is shown in the following chart.

- The death rate in LaSalle Parish is similar to Louisiana but worse than the service area and US.

Diabetes: Age-Adjusted Mortality Trends
(2007-2016 Annual Average Deaths per 100,000 Population)
Healthy People 2020 Target = 20.5 or Lower (Adjusted)



Sources: • CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics. Data extracted August 2018.


Notes: • US Department of Health and Human Services. Healthy People 2020. December 2010. <http://www.healthypeople.gov> [Objective D-3]
• Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10).
• Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population.
• The Healthy People 2020 target for Diabetes is adjusted to account for only diabetes mellitus coded deaths.

Prevalence of Diabetes

“Have you ever been told by a doctor, nurse, or other health professional that you have diabetes? (If female, add: not counting diabetes only occurring during pregnancy?)”

“Have you ever been told by a doctor, nurse, or other health professional that you have pre-diabetes or borderline diabetes? (If female, add: other than during pregnancy?)”

- Prevalence of diabetes in LaSalle Parish is less favorable than the statewide rate but similar to regional and national rates. The parish rate has fluctuated over time but is statistically similar to the baseline prevalence.




Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 140]
 • 2017 PRC National Health Survey, Professional Research Consultants, Inc.
 • Behavioral Risk Factor Surveillance System Survey Data. Atlanta, Georgia. United States Department of Health and Human Services, Centers for Disease Control and Prevention (CDC); 2016 LA data.

Notes: • Asked of all respondents.

- Seniors are much more likely than younger adults to be diabetic. There are no significant differences within the other demographic groups.

Prevalence of Diabetes (LaSalle Parish, 2018)




Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 140]
 Notes: • Asked of all respondents.
 • Income categories reflect respondent's household income as a ratio to the federal poverty level (FPL) for their household size. "Low Income" includes households with incomes up to 200% of the federal poverty level; "Mid/High Income" includes households with incomes at 200% or more of the federal poverty level.
 • Excludes gestational diabetes (occurring only during pregnancy).

Diabetes Testing

Adults who do not have diabetes: "Have you had a test for high blood sugar or diabetes within the past three years?"

- Testing levels in the parish are higher than found across the Rapides Foundation Service area and the US.

Have Had Blood Sugar Tested in the Past Three Years (Among Nondiabetics)



Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 39]
 • 2017 PRC National Health Survey, Professional Research Consultants, Inc.
 Notes: • Asked of respondents who have not been diagnosed with diabetes.

Key Informant Input: Diabetes

The following chart outlines key informants' perceptions of the severity of *Diabetes* as a problem in the community:

**Perceptions of Diabetes
as a Problem in the Community**
(Key Informants, 2018)

■ Major Problem ■ Moderate Problem ■ Minor Problem ■ No Problem At All



Sources: • PRC Online Key Informant Survey, Professional Research Consultants, Inc.
Notes: • Asked of all respondents.

Challenges

Among those rating this issue as a “major problem,” the biggest challenges for people with diabetes are seen as:

Obesity and Lifestyle

- Onset of diabetes is mainly attributed to obesity, diet, and genetics. Maintaining control of diabetes is a struggle with many individuals due to a lack of compliance with provider’s instructions. – Other Health Provider (LaSalle Parish)
- Food is the biggest challenge. The temptations are everywhere. LGH provides some diabetic in-services, but many patients don’t attend these. I think the challenge is getting information to diabetic patients that they could use to manage the disease. – Other Health Provider (LaSalle Parish)
- Weight gain. – Community Leader (LaSalle Parish)
- Diet and exercise. – Community Leader (LaSalle Parish)

Lack of Providers

- Barriers to physicians specializing in diabetes care. Barriers to access medical nutrition therapy in the community, thus lack of patient education to successfully manage this disease. – Community Leader (LaSalle Parish)

Access to Care/Services

- Getting the help and exercise that is needed to maintaining a healthy state of being. – Social Services Provider (LaSalle Parish)

Diagnosis/Treatment

- Lack of knowledge by health professionals as to symptoms and treatments. – Community Leader (LaSalle Parish)

Health Education and Awareness

- Education about diabetes, especially in the older generation. – Other Health Provider (LaSalle Parish)

Alzheimer's Disease

About Dementia

Dementia is the loss of cognitive functioning—thinking, remembering, and reasoning—to such an extent that it interferes with a person's daily life. Dementia is not a disease itself, but rather a set of symptoms. Memory loss is a common symptom of dementia, although memory loss by itself does not mean a person has dementia. Alzheimer's disease is the most common cause of dementia, accounting for the majority of all diagnosed cases.

Alzheimer's disease is the 6th leading cause of death among adults age 18 years and older. Estimates vary, but experts suggest that up to 5.1 million Americans age 65 years and older have Alzheimer's disease. These numbers are predicted to more than double by 2050 unless more effective ways to treat and prevent Alzheimer's disease are found.


- Healthy People 2020 (www.healthypeople.gov)

Age-Adjusted Alzheimer's Disease Deaths

Age-adjusted Alzheimer's disease mortality is outlined in the following chart.

- The parish death rate is lower than the regional rate, similar to the statewide rate, and higher than the national rate.

Alzheimer's Disease: Age-Adjusted Mortality Trends
(2012-2016 Annual Average Deaths per 100,000 Population)



Sources: • CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics. Data extracted August 2018.

Notes: • Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10).
• Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population.


Key Informant Input: Dementias, Including Alzheimer's Disease

The following chart outlines key informants' perceptions of the severity of *Dementias, Including Alzheimer's Disease* as a problem in the community:

Perceptions of Dementia/Alzheimer's Disease as a Problem in the Community

(Key Informants, 2018)

■ Major Problem ■ Moderate Problem ■ Minor Problem ■ No Problem At All



Sources: • PRC Online Key Informant Survey, Professional Research Consultants, Inc.
Notes: • Asked of all respondents.

Top Concerns

Among those rating this issue as a "major problem," reasons related to the following:

Prevalence/Incidence

Dementia seems to be more prevalent in a younger population now. – Other Health Provider (LaSalle Parish)

We have two nursing homes, and a big percentage have dementia. – Community Leader (LaSalle Parish)

Widespread disease affecting many families; no cure. – Community Leader (LaSalle Parish)

It seems everyone around me gets it. – Community Leader (LaSalle Parish)

Lack of Specialists

No known specialty programs for treating these patients, other than placement into nursing homes. Not sure that local physicians are looking for signs of dementia and Alzheimer's early. – Social Services Provider (LaSalle Parish)

Kidney Disease

About Kidney Disease

Chronic kidney disease and end-stage renal disease are significant public health problems in the United States and a major source of suffering and poor quality of life for those afflicted. They are responsible for premature death and exact a high economic price from both the private and public sectors. Nearly 25% of the Medicare budget is used to treat people with chronic kidney disease and end-stage renal disease.

Genetic determinants have a large influence on the development and progression of chronic kidney disease. It is not possible to alter a person's biology and genetic determinants; however, environmental influences and individual behaviors also have a significant influence on the development and progression of chronic kidney disease. As a result, some populations are disproportionately affected. Successful behavior modification is expected to have a positive influence on the disease.

Diabetes is the most common cause of kidney failure. The results of the Diabetes Prevention Program (DPP) funded by the National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK) show that moderate exercise, a healthier diet, and weight reduction can prevent development of type 2 diabetes in persons at risk.

- Healthy People 2020 (www.healthypeople.gov)

Age-Adjusted Kidney Disease Deaths

Age-adjusted kidney disease mortality is described in the following chart.

- The parish death rate is similar to the service area and Louisiana but much higher than the US.

Kidney Disease: Age-Adjusted Mortality Trends (2007-2016 Annual Average Deaths per 100,000 Population)




Sources: • CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics. Data extracted August 2018.

Notes: • Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10).
• Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population.

Key Informant Input: Kidney Disease

The following chart outlines key informants' perceptions of the severity of *Kidney Disease* as a problem in the community:

Perceptions of Kidney Disease as a Problem in the Community (Key Informants, 2018)



Sources: • PRC Online Key Informant Survey, Professional Research Consultants, Inc.
Notes: • Asked of all respondents.

Top Concerns

Among those rating this issue as a “major problem,” the following reason was given:

Access to Care/Services

We are approached oftentimes by patients or family members in need of dialysis. We have researched the possibility of providing this service at LaSalle General Hospital, but the expense of such a program is too great. – Other Health Provider (LaSalle Parish)

Potentially Disabling Conditions

About Arthritis, Osteoporosis, & Chronic Back Conditions

There are more than 100 types of arthritis. Arthritis commonly occurs with other chronic conditions, such as diabetes, heart disease, and obesity. Interventions to treat the pain and reduce the functional limitations from arthritis are important, and may also enable people with these other chronic conditions to be more physically active. Arthritis affects 1 in 5 adults and continues to be the most common cause of disability. It costs more than \$128 billion per year. All of the human and economic costs are projected to increase over time as the population ages. There are interventions that can reduce arthritis pain and functional limitations, but they remain underused. These include: increased physical activity; self-management education; and weight loss among overweight/obese adults.

Osteoporosis is a disease marked by reduced bone strength leading to an increased risk of fractures (broken bones). In the United States, an estimated 5.3 million people age 50 years and older have osteoporosis. Most of these people are women, but about 0.8 million are men. Just over 34 million more people, including 12 million men, have low bone mass, which puts them at increased risk for developing osteoporosis. Half of all women and as many as 1 in 4 men age 50 years and older will have an osteoporosis-related fracture in their lifetime.

Chronic back pain is common, costly, and potentially disabling. About 80% of Americans experience low back pain in their lifetime. It is estimated that each year:

- 15%-20% of the population develop protracted back pain.
- 2-8% have chronic back pain (pain that lasts more than 3 months).
- 3-4% of the population is temporarily disabled due to back pain.
- 1% of the working-age population is disabled completely and permanently as a result of low back pain.

Americans spend at least \$50 billion each year on low back pain. Low back pain is the:

- 2nd leading cause of lost work time (after the common cold).
- 3rd most common reason to undergo a surgical procedure.
- 5th most frequent cause of hospitalization.

Arthritis, osteoporosis, and chronic back conditions all have major effects on quality of life, the ability to work, and basic activities of daily living.

- Healthy People 2020 (www.healthypeople.gov)


Arthritis

“Would you please tell me if you have ever suffered from or been diagnosed with arthritis or rheumatism?”

See also *Overall Health Status: Activity Limitations* in the **General Health Status** section of this report.

- Prevalence of arthritis/rheumatism among those 50 and older in LaSalle Parish is higher than found across the nation.

Prevalence of Potentially Disabling Conditions




Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 141]
 • 2017 PRC National Health Survey, Professional Research Consultants, Inc.
 • US Department of Health and Human Services. Healthy People 2020. December 2010. <http://www.healthypeople.gov> [Objective AOCBC-10]
 Notes: • Reflects adults age 50+.

Key Informant Input: Arthritis, Osteoporosis, & Chronic Back Conditions

The following chart outlines key informants' perceptions of the severity of *Arthritis, Osteoporosis, & Chronic Back Conditions* as a problem in the community:

Perceptions of Arthritis/Osteoporosis/Back Conditions as a Problem in the Community (Key Informants, 2018)



Sources: • PRC Online Key Informant Survey, Professional Research Consultants, Inc.
 Notes: • Asked of all respondents.

Top Concerns

Among those rating this issue as a “major problem,” the following reason was given:

Diagnosis/Prevention

Osteoporosis isn't identified, even when family history indicates patients are high-risk. Back pain is such a prevalent patient complaint that it's often dismissed/not taken seriously, especially in a society where abuse of prescription painkillers is at epidemic heights. Ditto with arthritis: such the common ailment, yet quickly dismissed by health care providers as treatable by patients themselves with over-the-counter medications. While over-the-counter drugs may indeed help in all these situations, it is the initial diagnosis and primarily the initial lack of concern by the primary doctor that may lead patients to feel continued checkups are a waste of time and resources. – Community Leader (LaSalle Parish)

Vision & Hearing Impairment

Key Informant Input: Vision & Hearing

The following chart outlines key informants' perceptions of the severity of *Vision & Hearing* as a problem in the community:

Perceptions of Vision and Hearing as a Problem in the Community

(Key Informants, 2018)

■ Major Problem ■ Moderate Problem ■ Minor Problem ■ No Problem At All



Sources: • PRC Online Key Informant Survey, Professional Research Consultants, Inc.
Notes: • Asked of all respondents.

Infectious Disease

About Immunization & Infectious Diseases

The increase in life expectancy during the 20th century is largely due to improvements in child survival; this increase is associated with reductions in infectious disease mortality, due largely to immunization. However, infectious diseases remain a major cause of illness, disability, and death. Immunization recommendations in the United States currently target 17 vaccine-preventable diseases across the lifespan.

People in the US continue to get diseases that are vaccine-preventable. Viral hepatitis, influenza, and tuberculosis (TB) remain among the leading causes of illness and death across the nation and account for substantial spending on the related consequences of infection.

The infectious disease public health infrastructure, which carries out disease surveillance at the national, state, and local levels, is an essential tool in the fight against newly emerging and re-emerging infectious diseases. Other important defenses against infectious diseases include:

- Proper use of vaccines
- Antibiotics
- Screening and testing guidelines
- Scientific improvements in the diagnosis of infectious disease-related health concerns

Vaccines are among the most cost-effective clinical preventive services and are a core component of any preventive services package. Childhood immunization programs provide a very high return on investment. For example, for each birth cohort vaccinated with the routine immunization schedule, society:

- Saves 33,000 lives.
- Prevents 14 million cases of disease.
- Reduces direct healthcare costs by \$9.9 billion.
- Saves \$33.4 billion in indirect costs.
- Healthy People 2020 (www.healthypeople.gov)

Key Informant Input: Immunization & Infectious Diseases

The following chart outlines key informants' perceptions of the severity of *Immunization & Infectious Diseases* as a problem in the community:

Perceptions of Immunization and Infectious Diseases as a Problem in the Community

(Key Informants, 2018)

■ Major Problem ■ Moderate Problem ■ Minor Problem ■ No Problem At All



Sources: • PRC Online Key Informant Survey, Professional Research Consultants, Inc.
Notes: • Asked of all respondents.

Influenza & Pneumonia Vaccination

About Influenza & Pneumonia

Acute respiratory infections, including pneumonia and influenza, are the 8th leading cause of death in the nation, accounting for 56,000 deaths annually. Pneumonia mortality in children fell by 97% in the last century, but respiratory infectious diseases continue to be leading causes of pediatric hospitalization and outpatient visits in the US. On average, influenza leads to more than 200,000 hospitalizations and 36,000 deaths each year. The 2009 H1N1 influenza pandemic caused an estimated 270,000 hospitalizations and 12,270 deaths (1,270 of which were of people younger than age 18) between April 2009 and March 2010.

- Healthy People 2020 (www.healthypeople.gov)

Vaccinations


“During the past 12 months, have you had a flu shot?”

“A pneumonia shot or pneumococcal vaccine is usually given only once or twice in a person's lifetime and is different from the seasonal flu shot. Have you ever had a pneumonia shot?”

Columns in the following chart show these findings among those age 65+.

- The flu vaccination rate among LaSalle Parish seniors is higher than the statewide rate but comparable to service area and US rates. Despite a drop, the parish rate has returned to the baseline level.

Older Adults: Have Had a Flu Vaccination in the Past Year
(Among Adults Age 65+)
Healthy People 2020 Target = 70.0% or Higher




- Sources:
- 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 144]
 - 2017 PRC National Health Survey, Professional Research Consultants, Inc.
 - Behavioral Risk Factor Surveillance System Survey Data. Atlanta, Georgia. United States Department of Health and Human Services, Centers for Disease Control and Prevention (CDC): 2016 LA data.
 - US Department of Health and Human Services. Healthy People 2020. December 2010. <http://www.healthypeople.gov> [Objective IID-12.12]
- Notes:
- Reflects respondents 65 and older.
 - Includes Fluzone Intradermal as a form of vaccination.

- The pneumonia vaccination rate among parish seniors is statistically comparable to regional, state, and national rates. The parish prevalence has fluctuated dramatically since the 2002 rate.

Older Adults: Have Ever Had a Pneumonia Vaccine (Among Adults Age 65+)

Healthy People 2020 Target = 90.0% or Higher



- Sources:
- 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 146]
 - 2017 PRC National Health Survey, Professional Research Consultants, Inc.
 - Behavioral Risk Factor Surveillance System Survey Data. Atlanta, Georgia. United States Department of Health and Human Services, Centers for Disease Control and Prevention (CDC): 2016 LA data.
 - US Department of Health and Human Services. Healthy People 2020. December 2010. <http://www.healthypeople.gov> [Objectives IID-13.1, IID-13.2]
- Notes:
- Reflects respondents 65 and older.

HIV

About Human Immunodeficiency Virus (HIV)

The HIV epidemic in the United States continues to be a major public health crisis. An estimated 1.1 million Americans are living with HIV, and 1 in 5 people with HIV do not know they have it. HIV continues to spread, leading to about 56,000 new HIV infections each year.

HIV is a preventable disease, and effective HIV prevention interventions have been proven to reduce HIV transmission. People who get tested for HIV and learn that they are infected can make significant behavior changes to improve their health and reduce the risk of transmitting HIV to their sex or drug-using partners. More than 50% of new HIV infections occur as a result of the 21% of people who have HIV but do not know it.

In the era of increasingly effective treatments for HIV, people with HIV are living longer, healthier, and more productive lives. Deaths from HIV infection have greatly declined in the United States since the 1990s. As the number of people living with HIV grows, it will be more important than ever to increase national HIV prevention and healthcare programs.

There are gender, race, and ethnicity disparities in new HIV infections:

- Nearly 75% of new HIV infections occur in men.
- More than half occur in gay and bisexual men, regardless of race or ethnicity.
- 45% of new HIV infections occur in African Americans, 35% in whites, and 17% in Hispanics.

Improving access to quality healthcare for populations disproportionately affected by HIV, such as persons of color and gay and bisexual men, is a fundamental public health strategy for HIV prevention. People getting care for HIV can receive:

- Antiretroviral therapy
- Screening and treatment for other diseases (such as sexually transmitted infections)
- HIV prevention interventions
- Mental health services
- Other health services

As the number of people living with HIV increases and more people become aware of their HIV status, prevention strategies that are targeted specifically for HIV-infected people are becoming more important. Prevention work with people living with HIV focuses on:

- Linking to and staying in treatment.
- Increasing the availability of ongoing HIV prevention interventions.
- Providing prevention services for their partners.


Public perception in the US about the seriousness of the HIV epidemic has declined in recent years. There is evidence that risky behaviors may be increasing among uninfected people, especially gay and bisexual men. Ongoing media and social campaigns for the general public and HIV prevention interventions for uninfected persons who engage in risky behaviors are critical.

- Healthy People 2020 (www.healthypeople.gov)

HIV Prevalence

The following chart outlines prevalence (current cases, regardless of when they were diagnosed) of HIV per 100,000 population in the area.

- HIV prevalence in LaSalle Parish is considerably higher than found across the service area, state, and US.



Sources:


- Centers for Disease Control and Prevention, National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention.
- Retrieved August 2018 from Community Commons at <http://www.chna.org>.

Notes:

- This indicator is relevant because HIV is a life-threatening communicable disease that disproportionately affects minority populations and may also indicate the prevalence of unsafe sex practices.

Key Informant Input: HIV/AIDS

The following chart outlines key informants' perceptions of the severity of *HIV/AIDS* as a problem in the community:



Sources:

- PRC Online Key Informant Survey, Professional Research Consultants, Inc.

Notes:

- Asked of all respondents.

Sexually Transmitted Diseases

About Sexually Transmitted Diseases

STDs refer to more than 25 infectious organisms that are transmitted primarily through sexual activity. Despite their burdens, costs, and complications, and the fact that they are largely preventable, STDs remain a significant public health problem in the United States. This problem is largely unrecognized by the public, policymakers, and health care professionals. STDs cause many harmful, often irreversible, and costly clinical complications, such as: reproductive health problems; fetal and perinatal health problems; cancer; and facilitation of the sexual transmission of HIV infection.

Because many cases of STDs go undiagnosed—and some common viral infections, such as human papillomavirus (HPV) and genital herpes, are not reported to CDC at all—the reported cases of chlamydia, gonorrhea, and syphilis represent only a fraction of the true burden of STDs in the US. Untreated STDs can lead to serious long-term health consequences, especially for adolescent girls and young women. Several factors contribute to the spread of STDs.

Biological Factors. STDs are acquired during unprotected sex with an infected partner. Biological factors that affect the spread of STDs include:

- **Asymptomatic nature of STDs.** The majority of STDs either do not produce any symptoms or signs, or they produce symptoms so mild that they are unnoticed; consequently, many infected persons do not know that they need medical care.
- **Gender disparities.** Women suffer more frequent and more serious STD complications than men do. Among the most serious STD complications are pelvic inflammatory disease, ectopic pregnancy (pregnancy outside of the uterus), infertility, and chronic pelvic pain.
- **Age disparities.** Compared to older adults, sexually active adolescents ages 15 to 19 and young adults ages 20 to 24 are at higher risk for getting STDs.
- **Lag time between infection and complications.** Often, a long interval, sometimes years, occurs between acquiring an STD and recognizing a clinically significant health problem.

Social, Economic, and Behavioral Factors. The spread of STDs is directly affected by social, economic, and behavioral factors. Such factors may cause serious obstacles to STD prevention due to their influence on social and sexual networks, access to and provision of care, willingness to seek care, and social norms regarding sex and sexuality. Among certain vulnerable populations, historical experience with segregation and discrimination exacerbates these factors. Social, economic, and behavioral factors that affect the spread of STDs include: racial and ethnic disparities; poverty and marginalization; access to healthcare; substance abuse; sexuality and secrecy (stigma and discomfort discussing sex); and sexual networks (persons “linked” by sequential or concurrent sexual partners).

- Healthy People 2020 (www.healthypeople.gov)

Chlamydia & Gonorrhea


Chlamydia. Chlamydia is the most commonly reported STD in the United States; most people who have chlamydia are unaware, since the disease often has no symptoms.

Gonorrhea. Anyone who is sexually active can get gonorrhea. Gonorrhea can be cured with the right medication; left untreated, however, gonorrhea can cause serious health problems in both women and men.

The following chart outlines local incidence for these STDs.

- The parish rates for chlamydia and gonorrhea are considerably lower than the rates found across Louisiana and the US.

Chlamydia & Gonorrhea Incidence (Incidence Rate per 100,000 Population, 2014)




Sources: • Centers for Disease Control and Prevention, National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention.
 • Retrieved August 2018 from Community Commons at <http://www.chna.org>.
 Notes: • This indicator is relevant because it is a measure of poor health status and indicates the prevalence of unsafe sex practices.

Key Informant Input: Sexually Transmitted Diseases

The following chart outlines key informants' perceptions of the severity of *Sexually Transmitted Diseases* as a problem in the community:

Perceptions of Sexually Transmitted Diseases as a Problem in the Community (Key Informants, 2018)



Sources: • PRC Online Key Informant Survey, Professional Research Consultants, Inc.
 Notes: • Asked of all respondents.

Births

About Infant & Child Health

Improving the well-being of mothers, infants, and children is an important public health goal for the US. Their well-being determines the health of the next generation and can help predict future public health challenges for families, communities, and the healthcare system. The risk of maternal and infant mortality and pregnancy-related complications can be reduced by increasing access to quality preconception (before pregnancy) and inter-conception (between pregnancies) care. Moreover, healthy birth outcomes and early identification and treatment of health conditions among infants can prevent death or disability and enable children to reach their full potential. Many factors can affect pregnancy and childbirth, including pre-conception health status, age, access to appropriate healthcare, and poverty.

Infant and child health are similarly influenced by socio-demographic factors, such as family income, but are also linked to the physical and mental health of parents and caregivers. There are racial and ethnic disparities in mortality and morbidity for mothers and children, particularly for African Americans. These differences are likely the result of many factors, including social determinants (such as racial and ethnic disparities in infant mortality; family income; educational attainment among household members; and health insurance coverage) and physical determinants (i.e., the health, nutrition, and behaviors of the mother during pregnancy and early childhood).

- [Healthy People 2020 \(www.healthypeople.gov\)](http://www.healthypeople.gov)

Birth Outcomes & Risks

Low-Weight Births

Low birthweight babies, those who weigh less than 2,500 grams (5 pounds, 8 ounces) at birth, are much more prone to illness and neonatal death than are babies of normal birthweight. Largely a result of receiving poor or inadequate prenatal care, many low-weight births and the consequent health problems are preventable. Births of low-weight infants are described in the following chart.

- [The parish rate is lower than the statewide rate but nearly identical to the national rate.](#)

Low-Weight Births (Percent of Live Births)

Healthy People 2020 Target = 7.8% or Lower



Sources: • CDC WONDER Online Query System. Centers for Disease Control and Prevention, National Center for Health Statistics, Division of Vital Statistics. Data extracted August 2018.
 • US Department of Health and Human Services. Healthy People 2020. December 2010. <http://www.healthypeople.gov> [Objective MICH-8.1]
 Note: • This indicator reports the percentage of total births that are low birth weight (Under 2500g). This indicator is relevant because low birth weight infants are at high risk for health problems. This indicator can also highlight the existence of health disparities.

Infant Mortality


Infant mortality rates reflect deaths of children less than one year old per 1,000 live births. These rates are outlined in the following chart.

- The parish death rate is lower than regional and statewide rates but is nearly identical to the US rate.

Infant Mortality Rate

(2007-2016 Annual Average Infant Deaths per 1,000 Live Births)

Healthy People 2020 Target = 6.0 or Lower




Sources: • CDC WONDER Online Query System. Centers for Disease Control and Prevention, National Center for Health Statistics, Division of Vital Statistics. Data extracted August 2018.
 • Centers for Disease Control and Prevention, National Center for Health Statistics.
 • US Department of Health and Human Services. Healthy People 2020. December 2010. <http://www.healthypeople.gov> [Objective MICH-1.3]
 Notes: • Rates are three-year averages of deaths of children under 1 year old per 1,000 live births.

Key Informant Input: Infant & Child Health

The following chart outlines key informants' perceptions of the severity of *Infant & Child Health* as a problem in the community:

**Perceptions of Infant and Child Health
as a Problem in the Community**
(Key Informants, 2018)

■ Major Problem ■ Moderate Problem ■ Minor Problem ■ No Problem At All



Sources: • PRC Online Key Informant Survey, Professional Research Consultants, Inc.
Notes: • Asked of all respondents.

Family Planning

Births to Teen Mothers

About Teen Births

The negative outcomes associated with unintended pregnancies are compounded for adolescents. Teen mothers:


- Are less likely to graduate from high school or attain a GED by the time they reach age 30.
- Earn an average of approximately \$3,500 less per year, when compared with those who delay childbearing.
- Receive nearly twice as much Federal aid for nearly twice as long.

Similarly, early fatherhood is associated with lower educational attainment and lower income. Children of teen parents are more likely to have lower cognitive attainment and exhibit more behavior problems. Sons of teen mothers are more likely to be incarcerated, and daughters are more likely to become adolescent mothers.

- Healthy People 2020 (www.healthypeople.gov)

The following chart describes local teen births.

- The parish teen birth rate is statistically similar to Louisiana but much higher than the US. The decrease over time is not statistically significant.




Sources: • Centers for Disease Control and Prevention, National Vital Statistics System. Accessed using CDC WONDER.
• Retrieved from Community Commons at <http://www.chna.org>.

Notes: • This indicator reports the rate of total births to women under the age of 15–19 per 1,000 female population age 15–19. This indicator is relevant because in many cases, teen parents have unique social, economic, and health support services. Additionally, high rates of teen pregnancy may indicate the prevalence of unsafe sex practices.

- Note the difference in birth rate by race/ethnicity in LaSalle Parish when compared to the state and nation.

Teen Birth Rate

(Births to Women Age 15-19 Per 1,000 Female Population Age 15-19; LaSalle Parish by Race/Ethnicity, 2006-2012)




Sources: • Centers for Disease Control and Prevention, National Vital Statistics System. Accessed using CDC WONDER.
 • Retrieved from Community Commons at <http://www.chna.org>.
 Notes: • This indicator reports the rate of total births to women under the age of 15–19 per 1,000 female population age 15–19. This indicator is relevant because in many cases, teen parents have unique social, economic, and health support services. Additionally, high rates of teen pregnancy may indicate the prevalence of unsafe sex practices.

Key Informant Input: Family Planning

The following chart outlines key informants’ perceptions of the severity of *Family Planning* as a problem in the community:

Perceptions of Family Planning as a Problem in the Community (Key Informants, 2018)



Sources: • PRC Online Key Informant Survey, Professional Research Consultants, Inc.
 Notes: • Asked of all respondents.

Top Concerns

Among those rating this issue as a “major problem,” reasons related to the following:

Prevalence/Incidence

There are several teenage parents around the area. – Social Services Provider (LaSalle Parish)

Access to Care/Services

No facilities available. – Community Leader (LaSalle Parish)

Modifiable Health Risks

Nutrition, Physical Activity, & Weight

Nutrition

About Healthful Diet & Healthy Weight

Strong science exists supporting the health benefits of eating a healthful diet and maintaining a healthy body weight. Efforts to change diet and weight should address individual behaviors, as well as the policies and environments that support these behaviors in settings such as schools, worksites, healthcare organizations, and communities.

The goal of promoting healthful diets and healthy weight encompasses increasing household food security and eliminating hunger.

Americans with a healthful diet:

- Consume a variety of nutrient-dense foods within and across the food groups, especially whole grains, fruits, vegetables, low-fat or fat-free milk or milk products, and lean meats and other protein sources.
- Limit the intake of saturated and trans fats, cholesterol, added sugars, sodium (salt), and alcohol.
- Limit caloric intake to meet caloric needs.

Diet and body weight are related to health status. Good nutrition is important to the growth and development of children. A healthful diet also helps Americans reduce their risks for many health conditions, including: overweight and obesity; malnutrition; iron-deficiency anemia; heart disease; high blood pressure; dyslipidemia (poor lipid profiles); type 2 diabetes; osteoporosis; oral disease; constipation; diverticular disease; and some cancers.

Diet reflects the variety of foods and beverages consumed over time and in settings such as worksites, schools, restaurants, and the home. Interventions to support a healthier diet can help ensure that:

- Individuals have the knowledge and skills to make healthier choices.
- Healthier options are available and affordable.

Social Determinants of Diet. Demographic characteristics of those with a more healthful diet vary with the nutrient or food studied. However, most Americans need to improve some aspect of their diet.

Social factors thought to influence diet include:

- Knowledge and attitudes
- Skills
- Social support
- Societal and cultural norms
- Food and agricultural policies
- Food assistance programs
- Economic price systems

Physical Determinants of Diet. Access to and availability of healthier foods can help people follow healthful diets. For example, better access to retail venues that sell healthier options may have a positive impact on a person's diet; these venues may be less available in low-income or rural neighborhoods.

The places where people eat appear to influence their diet. For example, foods eaten away from home often have more calories and are of lower nutritional quality than foods prepared at home.

Marketing also influences people's—particularly children's—food choices.

- Healthy People 2020 (www.healthypeople.gov)

Daily Recommendation of Fruits/Vegetables

To measure fruit and vegetable consumption, survey respondents were asked multiple questions, specifically about the foods and drinks they consumed on the day prior to the interview.

“For the next questions, please think about the foods you ate yesterday. How many servings of fruit did you have yesterday?”

“How many servings of 100% fruit juice did you have yesterday?”


“How many servings of dark green or orange vegetables, such as carrots, broccoli, or sweet potatoes, did you have yesterday? (Examples of dark green vegetables are broccoli, spinach, collards, etc.) (Examples of orange vegetables are carrots and sweet potatoes, etc.)”

“How many servings of other vegetables did you have yesterday? (Examples are potatoes, corn, onions, peas, etc.)?”

The questions above are used to calculate daily fruit/vegetable consumption for respondents. The proportion reporting having 5 or more servings per day is shown here.

- The recommended consumption of fruits and vegetables in the parish is statistically similar to what is found across the country.
- There is no significant difference within the demographic groups.

Consume Five or More Servings of Fruits/Vegetables Per Day (LaSalle Parish, 2018)



Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 361]

Notes: • Asked of all respondents.

• Income categories reflect respondent's household income as a ratio to the federal poverty level (FPL) for their household size. "Low Income" includes households with incomes up to 200% of the federal poverty level; "Mid/High Income" includes households with incomes at 200% or more of the federal poverty level.


• For this issue, respondents were asked to recall their food intake on the previous day.

Access to Fresh Produce

“How difficult is it for you to buy fresh produce like fruits and vegetables at a price you can afford — would you say: very difficult, somewhat difficult, not too difficult, or not at all difficult?”

- The parish response is identical to the RFSA response and statistically unchanged over time.

Find It “Very” or “Somewhat” Difficult to Buy Affordable Fresh Produce




Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 321]
 Notes: • Asked of all respondents.

A food desert is defined as a low-income area where a significant number or share of residents is far from a supermarket, where "far" is more than 1 mile in urban areas and more than 10 miles in rural areas. This related chart is based on US Department of Agriculture data.

- Similar proportions of low food access are found across the parish, service area, and state.

Population With Low Food Access

(Percent of Population That Is Far From a Supermarket or Large Grocery Store, 2015)



Sources: • US Department of Agriculture, Economic Research Service, USDA - Food Access Research Atlas (FARA).
 • Retrieved August 2018 from Community Commons at <http://www.chna.org>.
 Notes: • This indicator reports the percentage of the population living in census tracts designated as food deserts. A food desert is defined as low-income areas where a significant number or share of residents is far from a supermarket, where "far" is more than 1 mile in urban areas and more than 10 miles in rural areas. This indicator is relevant because it highlights populations and geographies facing food insecurity.

Physical Activity

About Physical Activity

Regular physical activity can improve the health and quality of life of Americans of all ages, regardless of the presence of a chronic disease or disability. Among adults, physical activity can lower the risk of: early death; coronary heart disease; stroke; high blood pressure; type 2 diabetes; breast and colon cancer; falls; and depression. Among children and adolescents, physical activity can: improve bone health; improve cardiorespiratory and muscular fitness; decrease levels of body fat; and reduce symptoms of depression. For people who are inactive, even small increases in physical activity are associated with health benefits.

Personal, social, economic, and environmental factors all play a role in physical activity levels among youth, adults, and older adults. Understanding the barriers to and facilitators of physical activity is important to ensure the effectiveness of interventions and other actions to improve levels of physical activity.

Factors **positively** associated with adult physical activity include: postsecondary education; higher income; enjoyment of exercise; expectation of benefits; belief in ability to exercise (self-efficacy); history of activity in adulthood; social support from peers, family, or spouse; access to and satisfaction with facilities; enjoyable scenery; and safe neighborhoods.

Factors **negatively** associated with adult physical activity include: advancing age; low income; lack of time; low motivation; rural residency; perception of great effort needed for exercise; overweight or obesity; perception of poor health; and being disabled. Older adults may have additional factors that keep them from being physically active, including lack of social support, lack of transportation to facilities, fear of injury, and cost of programs.

Among children ages 4 to 12, the following factors have a positive association with physical activity: gender (boys); belief in ability to be active (self-efficacy); and parental support.

Among adolescents ages 13 to 18, the following factors have a positive association with physical activity: parental education; gender (boys); personal goals; physical education/school sports; belief in ability to be active (self-efficacy); and support of friends and family.

Environmental influences positively associated with physical activity among children and adolescents include:

- Presence of sidewalks
- Having a destination/walking to a particular place
- Access to public transportation
- Low traffic density
- Access to neighborhood or school play area and/or recreational equipment

People with disabilities may be less likely to participate in physical activity due to physical, emotional, and psychological barriers. Barriers may include the inaccessibility of facilities and the lack of staff trained in working with people with disabilities.

- Healthy People 2020 (www.healthypeople.gov)

Leisure-Time Physical Activity


Leisure-time physical activity includes any physical activities or exercises (such as running, calisthenics, golf, gardening, walking, etc.) which take place outside of one's line of work.

“During the past month, other than your regular job, did you participate in any physical activities or exercises, such as running, calisthenics, golf, gardening, or walking for exercise?”

- Lack of leisure-time physical activity in the parish is statistically similar to the populations shown in the following chart. The parish rate is comparable to the baseline rate.

No Leisure-Time Physical Activity in the Past Month

Healthy People 2020 Target = 32.6% or Lower



Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 106]
 • Behavioral Risk Factor Surveillance System Survey Data. Atlanta, Georgia. United States Department of Health and Human Services, Centers for Disease Control and Prevention (CDC); 2016 LA data.
 • 2017 PRC National Health Survey, Professional Research Consultants, Inc.
 • US Department of Health and Human Services. Healthy People 2020. December 2010. <http://www.healthypeople.gov> [Objective PA-1]

Notes: • Asked of all respondents.

Recommended Levels of Physical Activity

Adults should do 2 hours and 30 minutes a week of moderate-intensity (such as walking), or 1 hour and 15 minutes (75 minutes) a week of vigorous-intensity aerobic physical activity (such as jogging), or an equivalent combination of moderate- and vigorous-intensity aerobic physical activity. The guidelines also recommend that adults do muscle-strengthening activities, such as push-ups, sit-ups, or activities using resistance bands or weights. These activities should involve all major muscle groups and be done on two or more days per week.

The report finds that nationwide nearly 50 percent of adults are getting the recommended amounts of aerobic activity and about 30 percent are engaging in the recommended muscle-strengthening activity.

- 2013 Physical Activity Guidelines for Americans, US Department of Health and Human Services. www.cdc.gov/physicalactivity

Meeting Physical Activity Recommendations

To measure physical activity frequency, duration and intensity, respondents were asked:

“During the past month, what type of physical activity or exercise did you spend the most time doing?”

“And during the past month, how many times per week or per month did you take part in this activity?”

“And when you took part in this activity, for how many minutes or hours did you usually keep at it?”

Respondents could answer the above series for up to two types of physical activity. The specific activities identified (e.g., jogging, basketball, treadmill, etc.) determined the intensity values assigned to that respondent when calculating total aerobic physical activity hours/minutes.

Respondents were also asked about strengthening exercises:


“During the past month, how many times per week or per month did you do physical activities or exercises to strengthen your muscles? Do not count aerobic activities like walking, running, or bicycling. Please include activities using your own body weight, such as yoga, sit-ups, or push-ups, and those using weight machines, free weights, or elastic bands.”

“Meeting physical activity recommendations” includes adequate levels of both aerobic and strengthening activity:

- Aerobic activity is at least 150 minutes per week of light to moderate activity, 75 minutes per week of vigorous physical activity, or an equivalent combination of both;
- Strengthening activity is at least 2 sessions per week of exercise designed to strengthen muscles.
- The proportion of those in the parish who meet both recommendations is statistically comparable to the national figure.
- Within the demographic groups, there are no statistically significant differences.

Meets Physical Activity Recommendations (LaSalle Parish, 2018)

Healthy People 2020 Target = 20.1% or Higher



Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 152]
 • US Department of Health and Human Services. Healthy People 2020. December 2010. <http://www.healthypeople.gov> [Objective PA-2.4]


Notes: • Asked of all respondents.
 • Income categories reflect respondent's household income as a ratio to the federal poverty level (FPL) for their household size. "Low Income" includes households with incomes up to 200% of the federal poverty level; "Mid/High Income" includes households with incomes at 200% or more of the federal poverty level.
 • Meeting both guidelines is defined as the number of persons age 18+ who report light or moderate aerobic activity for at least 150 minutes per week or who report vigorous physical activity 75 minutes per week or an equivalent combination of moderate and vigorous-intensity activity and report doing physical activities specifically designed to strengthen muscles at least twice per week.

Walking

“How many days per week or per month do you walk for more than 10 minutes at a time?”

- The parish response is comparable to the Rapides Foundation Service Area response and the 2010 response, though marking an increase from 2013 survey results.

Walk for More Than 10 Minutes at a Time at Least Five Times per Week




Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 366]
 Notes: • Asked of all respondents.

Community Opportunities for Physical Activity

“How would you rate the availability of opportunities to participate in physical activity in your community? Would you say: excellent, very good, good, fair, or poor?”

- “Fair” or “poor” responses in LaSalle Parish are similar to the Rapides Foundation Service Area. The parish rate is lower than the 2010 rate.

“Fair” or “Poor” Evaluations of the Availability of Opportunities to Participate in Physical Activity in the Community



Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 327]
 Notes: • Asked of all respondents.

Weight Status

About Overweight & Obesity

Because weight is influenced by energy (calories) consumed and expended, interventions to improve weight can support changes in diet or physical activity. They can help change individuals' knowledge and skills, reduce exposure to foods low in nutritional value and high in calories, or increase opportunities for physical activity. Interventions can help prevent unhealthy weight gain or facilitate weight loss among obese people. They can be delivered in multiple settings, including healthcare settings, worksites, or schools.

The social and physical factors affecting diet and physical activity (see Physical Activity topic area) may also have an impact on weight. Obesity is a problem throughout the population. However, among adults, the prevalence is highest for middle-aged people and for non-Hispanic black and Mexican American women. Among children and adolescents, the prevalence of obesity is highest among older and Mexican American children and non-Hispanic black girls. The association of income with obesity varies by age, gender, and race/ethnicity.

- Healthy People 2020 (www.healthypeople.gov)

Body Mass Index (BMI), which describes relative weight for height, is significantly correlated with total body fat content. The BMI should be used to assess overweight and obesity and to monitor changes in body weight. In addition, measurements of body weight alone can be used to determine efficacy of weight loss therapy. BMI is calculated as weight (kg)/height squared (m²). To estimate BMI using pounds and inches, use: [weight (pounds)/height squared (inches²)] x 703.

In this report, overweight is defined as a BMI of 25.0 to 29.9 kg/m² and obesity as a BMI ≥30 kg/m². The rationale behind these definitions is based on epidemiological data that show increases in mortality with BMIs above 25 kg/m². The increase in mortality, however, tends to be modest until a BMI of 30 kg/m² is reached. For persons with a BMI ≥30 kg/m², mortality rates from all causes, and especially from cardiovascular disease, are generally increased by 50 to 100 percent above that of persons with BMIs in the range of 20 to 25 kg/m².

- Clinical Guidelines on the Identification, Evaluation, and Treatment of Overweight and Obesity in Adults: The Evidence Report. National Institutes of Health. National Heart, Lung, and Blood Institute in Cooperation With The National Institute of Diabetes and Digestive and Kidney Diseases. September 1998.

Classification of Overweight and Obesity by BMI	BMI (kg/m ²)
Underweight	<18.5
Healthy Weight	18.5 – 24.9
Overweight, not Obese	25.0 – 29.9
Obese	≥30.0

Source: Clinical Guidelines on the Identification, Evaluation, and Treatment of Overweight and Obesity in Adults: The Evidence Report. National Institutes of Health. National Heart, Lung, and Blood Institute in Cooperation With The National Institute of Diabetes and Digestive and Kidney Diseases. September 1998.


“About how much do you weigh without shoes?”

“About how tall are you without shoes?”

“Are you now trying to lose weight?”

Reported height and weight were used to calculate a Body Mass Index or BMI value (described above) for each respondent. This calculation allows us to examine the proportion of the population who is at a healthy weight, or who is overweight or obese (see table above).


- The overweight prevalence in the parish is nearly identical to the service area but is less favorable than the state and nation. The parish rate is considerably higher than the 2002 baseline rate.



Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Items 154, 367]
 • 2017 PRC National Health Survey, Professional Research Consultants, Inc.
 • Behavioral Risk Factor Surveillance System Survey Data. Atlanta, Georgia. United States Department of Health and Human Services, Centers for Disease Control and Prevention (CDC): 2016 LA data.

Notes: • Based on reported heights and weights, asked of all respondents.
 • The definition of overweight is having a body mass index (BMI), a ratio of weight to height (kilograms divided by meters squared), greater than or equal to 25.0, regardless of gender. The definition for obesity is a BMI greater than or equal to 30.0.


- Prevalence of obesity in LaSalle Parish is statistically comparable to the populations shown below. The increase over time is not statistically significant.



Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 176]
 • 2017 PRC National Health Survey, Professional Research Consultants, Inc.
 • US Department of Health and Human Services. Healthy People 2020. December 2010. <http://www.healthypeople.gov> [Objective NWS-9]
 • Behavioral Risk Factor Surveillance System Survey Data. Atlanta, Georgia. United States Department of Health and Human Services, Centers for Disease Control and Prevention (CDC): 2016 LA data.


Notes: • Based on reported heights and weights, asked of all respondents.
 • The definition of obesity is having a body mass index (BMI), a ratio of weight to height (kilograms divided by meters squared), greater than or equal to 30.0, regardless of gender.

- Obesity prevalence is statistically comparable by demographics in LaSalle Parish.



Key Informant Input: Nutrition, Physical Activity, & Weight

The following chart outlines key informants' perceptions of the severity of *Nutrition, Physical Activity, & Weight* as a problem in the community:



Top Concerns

Among those rating this issue as a "major problem," reasons related to the following:

Access to Healthy Foods/Nutrition

Healthy foods cost more, and lack of exercise. – Community Leader (LaSalle Parish)

Health Education and Awareness

Education, time constraints for individuals, access to nutrition education. – Community Leader (LaSalle Parish)

Lack of Programs

No programs to really encourage handling these problems. No known place where individual needs for diet and activity are addressed. – Social Services Provider (LaSalle Parish)

Lifestyle

Biggest challenges are not just education on proper diet, but change of eating habits. Lack of physical activity and diet, of course, attribute to excess weight or obesity. Personal responsibility must be addressed. – Other Health Provider (LaSalle Parish)

Obesity

Rampant obesity and poor eating habits. – Community Leader (LaSalle Parish)

Substance Abuse

About Substance Abuse

Substance abuse has a major impact on individuals, families, and communities. The effects of substance abuse are cumulative, significantly contributing to costly social, physical, mental, and public health problems. These problems include:

- Teenage pregnancy
- Human immunodeficiency virus/acquired immunodeficiency syndrome (HIV/AIDS)
- Other sexually transmitted diseases (STDs)
- Domestic violence
- Child abuse
- Motor vehicle crashes
- Physical fights
- Crime
- Homicide
- Suicide

Substance abuse refers to a set of related conditions associated with the consumption of mind- and behavior-altering substances that have negative behavioral and health outcomes. Social attitudes and political and legal responses to the consumption of alcohol and illicit drugs make substance abuse one of the most complex public health issues. In addition to the considerable health implications, substance abuse has been a flash-point in the criminal justice system and a major focal point in discussions about social values: people argue over whether substance abuse is a disease with genetic and biological foundations or a matter of personal choice.

Advances in research have led to the development of evidence-based strategies to effectively address substance abuse. Improvements in brain-imaging technologies and the development of medications that assist in treatment have gradually shifted the research community's perspective on substance abuse. There is now a deeper understanding of substance abuse as a disorder that develops in adolescence and, for some individuals, will develop into a chronic illness that will require lifelong monitoring and care.

Improved evaluation of community-level prevention has enhanced researchers' understanding of environmental and social factors that contribute to the initiation and abuse of alcohol and illicit drugs, leading to a more sophisticated understanding of how to implement evidence-based strategies in specific social and cultural settings.

A stronger emphasis on evaluation has expanded evidence-based practices for drug and alcohol treatment. Improvements have focused on the development of better clinical interventions through research and increasing the skills and qualifications of treatment providers.


- Healthy People 2020 (www.healthypeople.gov)

Related Age-Adjusted Mortality

Unintentional Drug-Related Deaths. Unintentional drug-related deaths include all deaths, other than suicide, for which drugs are the underlying cause. A "drug" includes illicit or street drugs (e.g., heroin and cocaine), as well as legal prescription and over-the-counter drugs; alcohol is not included. The following chart outlines local age-adjusted mortality for unintentional drug-related deaths.

- The parish death rate is higher than is found across the region, state, and nation.

**Unintentional Drug-Related Deaths:
Age-Adjusted Mortality Trends**
(2007-2016 Annual Average Deaths per 100,000 Population)
Healthy People 2020 Target = 11.3 or Lower



Sources: • CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics. Data extracted August 2018.
• UD Department of Health and Human Services. Healthy People 2020. December 2010. <http://www.healthypeople.gov> [Objective SA-12].

Notes: • Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10).
• Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population.

Alcohol Use

Excessive Drinkers. Excessive drinking reflects the number of adults (age 18+) who drank more than two drinks per day on average (for men) or more than one drink per day on average (for women), or who drank 5 or more drinks during a single occasion (for men) or 4 or more drinks during a single occasion (for women) during the past 30 days.

“During the past 30 days, on how many days did you have at least one drink of any alcoholic beverage such as beer, wine, a malt beverage, or liquor?”


“On the day(s) when you drank, about how many drinks did you have on the average?”

“Considering all types of alcoholic beverages, how many times during the past 30 days did you have 5 (if male)/4 (if female) or more drinks on an occasion?”

- The rate of excessive drinking in LaSalle Parish is considerably lower than found nationwide.
- Men and adults younger than 65 are more likely to engage in excessive drinking. There is no significant variance by income.

Excessive Drinkers (LaSalle Parish, 2018)

Healthy People 2020 Target = 25.4% or Lower




- Sources:
- 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 168]
 - US Department of Health and Human Services. Healthy People 2020. December 2010. <http://www.healthypeople.gov> [Objective SA-15]
- Notes:
- Asked of all respondents.
 - Income categories reflect respondent's household income as a ratio to the federal poverty level (FPL) for their household size. "Low Income" includes households with incomes up to 200% of the federal poverty level; "Mid/High Income" includes households with incomes at 200% or more of the federal poverty level.
 - Excessive drinking reflects the number of persons aged 18 years and over who drank more than two drinks per day on average (for men) or more than one drink per day on average (for women) OR who drank 5 or more drinks during a single occasion (for men) or 4 or more drinks during a single occasion (for women) during the past 30 days.

Drinking & Driving. As a self-reported measure – and because this indicator reflects potentially illegal behavior – it is reasonable to expect that it might be underreported, and that the actual incidence of drinking and driving in the community is likely higher.

“During the past 30 days, how many times have you driven when you've had perhaps too much to drink?”

- The parish drinking and driving rate is statistically similar to the populations shown below. It is significantly higher than the baseline rate.

Have Driven in the Past Month After Perhaps Having Too Much to Drink



Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 66]
 • Behavioral Risk Factor Surveillance System Survey Data. Atlanta, Georgia. United States Department of Health and Human Services, Centers for Disease Control and Prevention (CDC): 2016 LA data.
 • 2017 PRC National Health Survey, Professional Research Consultants, Inc.
 Notes: • Asked of all respondents.


Illicit Drug Use

“During the past 30 days, have you used an illegal drug or taken a prescription drug that was not prescribed to you?”

- The parish rate of 0.0% is well below the regional and national rates and has not varied statistically over time.

Illicit Drug Use in the Past Month

Healthy People 2020 Target = 7.1% or Lower




Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 67]
 • 2017 PRC National Health Survey, Professional Research Consultants, Inc.
 • US Department of Health and Human Services. Healthy People 2020. December 2010. <http://www.healthypeople.gov> [Objective SA-13.3]
 Notes: • Asked of all respondents.

Use of Opiates/Opioids

“Opiates or opioids are drugs that doctors prescribe to treat pain. Examples of prescription opiates include morphine, codeine, hydrocodone, oxycodone, methadone, and fentanyl. In the past year, have you used any of these prescription opiates, whether or not a doctor had prescribed them to you?”

- Prescription opioid use is comparable across the parish and region.

Used Prescription Opioids or Opiates in the Past Year (Whether Prescribed or Not)
(LaSalle Parish, 2018)




Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 314]
 Notes: • Asked of all respondents.
 • Examples of prescription opiates include morphine, codeine, hydrocodone, oxycodone, methadone, and fentanyl.

Alcohol & Drug Treatment

“Have you ever sought professional help for an alcohol or drug-related problem?”

- The parish rate is similar to service area and US rates and statistically unchanged over time.

Have Ever Sought Professional Help for an Alcohol/Drug-Related Problem



Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 68]
 • 2017 PRC National Health Survey, Professional Research Consultants, Inc.
 Notes: • Asked of all respondents.

Key Informant Input: Substance Abuse

The following chart outlines key informants' perceptions of the severity of *Substance Abuse* as a problem in the community:

Perceptions of Substance Abuse as a Problem in the Community

(Key Informants, 2018)

■ Major Problem ■ Moderate Problem ■ Minor Problem ■ No Problem At All



Sources: • PRC Online Key Informant Survey, Professional Research Consultants, Inc.
Notes: • Asked of all respondents.

Barriers to Treatment

Among those rating this issue as a “major problem,” the greatest barriers to accessing substance abuse treatment are viewed as:

Access to Resources

A hospital provides a 3-day treatment facility for patients. No providers in the parish for patients to follow-up or to devise an outpatient program. Monetary barriers. Those who have insurance can only use this for so long, and insurance will deny payments. – Community Leader (LaSalle Parish)

Lack of substance abuse treatment and lack of insurance coverage. – Other Health Provider (LaSalle Parish)

Lack of Facilities/Providers

No programs or facilities for treatment or information on substance abuse. – Social Services Provider (LaSalle Parish)

Denial/Stigma

Ignorance by many people that there is a problem. – Community Leader (LaSalle Parish)

Lack of Funding

Money and local treatment centers. – Community Leader (LaSalle Parish)

Tobacco Use

About Tobacco Use

Tobacco use is the single most preventable cause of death and disease in the United States. Scientific knowledge about the health effects of tobacco use has increased greatly since the first Surgeon General's report on tobacco was released in 1964.

Tobacco use causes:

- Cancer
- Heart disease
- Lung diseases (including emphysema, bronchitis, and chronic airway obstruction)
- Premature birth, low birth weight, stillbirth, and infant death

There is no risk-free level of exposure to secondhand smoke. Secondhand smoke causes heart disease and lung cancer in adults and a number of health problems in infants and children, including: severe asthma attacks; respiratory infections; ear infections; and sudden infant death syndrome (SIDS).


Smokeless tobacco causes a number of serious oral health problems, including cancer of the mouth and gums, periodontitis, and tooth loss. Cigar use causes cancer of the larynx, mouth, esophagus, and lung.

- Healthy People 2020 (www.healthypeople.gov)

Cigarette Smoking

“Do you now smoke cigarettes every day, some days, or not at all?”

- The proportion of smokers in LaSalle Parish is lower than the service area and the state but similar to the US. Cigarette smoking in LaSalle Parish has decreased significantly since 2002.




- Sources:
- 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 181]
 - 2017 PRC National Health Survey, Professional Research Consultants, Inc.
 - Behavioral Risk Factor Surveillance System Survey Data. Atlanta, Georgia. United States Department of Health and Human Services, Centers for Disease Control and Prevention (CDC): 2016 LA data.
 - US Department of Health and Human Services. Healthy People 2020. December 2010. <http://www.healthypeople.gov> [Objective TU-1.1]
- Notes:
- Asked of all respondents.
 - Includes regular and occasional smokers (those who smoke cigarettes every day or on some days).

- Men are much more likely than women to smoke cigarettes. There is no significant difference within the other groups.

Current Smokers (LaSalle Parish, 2018)

Healthy People 2020 Target = 12.0% or Lower



Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 159]

• US Department of Health and Human Services. Healthy People 2020. December 2010. <http://www.healthypeople.gov> [Objective TU-1.1]

Notes: • Asked of all respondents.

• Income categories reflect respondent's household income as a ratio to the federal poverty level (FPL) for their household size. "Low Income" includes households with incomes up to 200% of the federal poverty level; "Mid/High Income" includes households with incomes at 200% or more of the federal poverty level.

• Includes regular and occasion smokers (every day and some days).

Smoking Cessation

About Reducing Tobacco Use

Preventing tobacco use and helping tobacco users quit can improve the health and quality of life for Americans of all ages. People who stop smoking greatly reduce their risk of disease and premature death. Benefits are greater for people who stop at earlier ages, but quitting tobacco use is beneficial at any age.

Many factors influence tobacco use, disease, and mortality. Risk factors include race/ethnicity, age, education, and socioeconomic status. Significant disparities in tobacco use exist geographically; such disparities typically result from differences among states in smoke-free protections, tobacco prices, and program funding for tobacco prevention.


- Healthy People 2020 (www.healthypeople.gov)

“Are you aware of any services, programs or classes in your area to help smokers quit smoking?”

(Asked of all respondents.)

- Awareness of smoking cessation programs is considerably lower in LaSalle Parish than in the Rapides Foundation Service Area. The parish rate is statistically similar to 2010 survey results.

Aware of Services, Programs, or Classes to Help Smokers Quit Smoking



Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 310]
 Notes: • Asked of all respondents.


Secondhand Smoke

“In the past 30 days, has anyone, including yourself, smoked cigarettes, cigars or pipes anywhere in your home on an average of four or more days per week?”

The following chart details these responses among the total sample of respondents.

- The parish household smoking rate is statistically similar to the service area and the US. Though fluctuating over time, the parish rate is similar to the 2005 study.

Member of Household Smokes at Home




Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Items 58, 184]
 • 2017 PRC National Health Survey, Professional Research Consultants, Inc.
 Notes: • Asked of all respondents.
 • “Smokes at home” refers to someone smoking cigarettes, cigars, or a pipe in the home an average of four or more times per week in the past month.

Use of Vaping Products

“The next questions are about electronic vaping products, such as electronic cigarettes, also known as e-cigarettes. These are battery-operated devices that simulate traditional cigarette smoking, but do not involve the burning of tobacco. The cartridge or liquid "e-juice" used in these devices produces vapor and comes in a variety of flavors. Have you ever used an electronic vaping product, such as an e-cigarette, even just one time in your entire life?”

“Do you now use electronic vaping products, such as e-cigarettes, "every day," "some days," or "not at all"?”

Use Vaping Products
(LaSalle Parish, 2018)




Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 54]
Notes: • Asked of all respondents.

Smokeless Tobacco

“Do you currently use chewing tobacco, snuff, or snus every day, some days, or not at all?”

- Smokeless tobacco use in LaSalle Parish is comparable to the service area but higher than the state and nation. The parish rate has increased considerably over time.

Use of Smokeless Tobacco Healthy People 2020 Target = 0.3% or Lower




Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 312]
 • 2017 PRC National Health Survey, Professional Research Consultants, Inc.
 • Behavioral Risk Factor Surveillance System Survey Data. Atlanta, Georgia. United States Department of Health and Human Services, Centers for Disease Control and Prevention (CDC): 2016 Louisiana data.
 • US Department of Health and Human Services. Healthy People 2020. December 2010. <http://www.healthypeople.gov> [Objective TU-1.2]
 Notes: • Asked of all respondents.
 • Includes chewing tobacco, snuff, or snus.

Key Informant Input: Tobacco Use

The following chart outlines key informants' perceptions of the severity of *Tobacco Use* as a problem in the community:

Perceptions of Tobacco Use as a Problem in the Community (Key Informants, 2018)



Sources: • PRC Online Key Informant Survey, Professional Research Consultants, Inc.
 Notes: • Asked of all respondents.

Top Concerns

Among those rating this issue as a “major problem,” reasons related to the following:

Prevalence/Incidence

There appears to be many smokers in the area. There does not appear to be any programs to inform, treat or discourage use of tobacco. – Social Services Provider (LaSalle Parish)

A lot of young men are dipping or chewing tobacco. – Community Leader (LaSalle Parish)

Culture/Social Norms

Cultural. – Other Health Provider (LaSalle Parish)

Teens/Young Adult Usage

Students as young as elementary school are regularly using tobacco products. It is a major concern at the high school level. Most adults and children in the community do not associate smokeless tobacco products with health issues. – Community Leader (LaSalle Parish)

Easily Accessible

Easy access to them, and it is more accepted in the community. More high school kids are starting to smoke. – Community Leader (LaSalle Parish)

Policies

Not many areas within community are deemed smoke free. No way to enforce these areas. – Community Leader (LaSalle Parish)

Access to Health Services


Lack of Health Insurance Coverage (Age 18 to 64)

Survey respondents were asked a series of questions to determine their healthcare insurance coverage, if any, from either private or government-sponsored sources. Here, lack of health insurance coverage reflects respondents age 18 to 64 (thus excluding the Medicare population), who have no type of insurance coverage for healthcare services – neither private insurance nor government-sponsored plans (e.g., Medicaid).

“Do you have any government-assisted healthcare coverage, such as Medicare, Medicaid (or another state-sponsored program), or VA/military benefits?”


“Do you currently have: health insurance you get through your own or someone else's employer or union; health insurance you purchase yourself; or, you do not have health insurance and pay for health care entirely on your own?”

- The proportion of uninsured adults in LaSalle Parish is similar to the service area and nation but better than Louisiana. The parish rate has improved over time.



- Lack of coverage is higher among parish men.

Lack of Healthcare Insurance Coverage (Among Adults Age 18-64; LaSalle Parish, 2018) Healthy People 2020 Target = 0.0% (Universal Coverage)



- Sources:
- 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 169]
 - US Department of Health and Human Services. Healthy People 2020. December 2010. <http://www.healthypeople.gov> [Objective AHS-1]
- Notes:
- Asked of all respondents under the age of 65.
 - Income categories reflect respondent's household income as a ratio to the federal poverty level (FPL) for their household size. "Low Income" includes households with incomes up to 200% of the federal poverty level; "Mid/High Income" includes households with incomes at 200% or more of the federal poverty level.

Difficulties Accessing Healthcare

About Access to Healthcare

Access to comprehensive, quality health care services is important for the achievement of health equity and for increasing the quality of a healthy life for everyone. It impacts: overall physical, social, and mental health status; prevention of disease and disability; detection and treatment of health conditions; quality of life; preventable death; and life expectancy.

Access to health services means the timely use of personal health services to achieve the best health outcomes. It requires three distinct steps: 1) Gaining entry into the health care system; 2) Accessing a health care location where needed services are provided; and 3) Finding a health care provider with whom the patient can communicate and trust.

- Healthy People 2020 (www.healthypeople.gov)

Barriers to Healthcare Access

To better understand healthcare access barriers, survey participants were asked whether any of the following barriers to access prevented them from seeing a physician or obtaining a needed prescription in the past year.


“Was there a time in the past 12 months when...

- ... you needed medical care, but had **difficulty finding a doctor?**”
- ... you had difficulty getting an **appointment** to see a doctor?”
- ... you needed to see a doctor, but could not because of the **cost?**”
- ... a **lack of transportation** made it difficult or prevented you from seeing a doctor or making a medical appointment?”
- ... you were not able to see a doctor because the **office hours were not convenient?**”
- ... you needed a **prescription medicine**, but did not get it because you could not afford it?”

The percentages shown in the following chart reflect the total population, regardless of whether medical care was needed or sought.

- All barriers to access shown in the following chart are statistically similar between the parish and US.


Barriers to Access Have Prevented Medical Care in the Past Year



The following charts reflect the composite percentage of the total population experiencing problems accessing healthcare in the past year (indicating one or more of the aforementioned barriers or any other problem not specifically asked), again regardless of whether they needed or sought care.


- The parish composite is statistically comparable to the regional and national composites. The parish rate has not varied statistically since 2002.

Experienced Difficulties or Delays of Some Kind in Receiving Needed Healthcare in the Past Year



- Those younger than 65 were much more likely than seniors to have difficulties accessing healthcare in the past year. There is no significant difference within the other groups.

Experienced Difficulties or Delays of Some Kind in Receiving Needed Healthcare in the Past Year (LaSalle Parish, 2018)




Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 171]
 Notes: • Asked of all respondents.
 • Represents the percentage of respondents experiencing one or more barriers to accessing healthcare in the past 12 months.
 • Income categories reflect respondent's household income as a ratio to the federal poverty level (FPL) for their household size. "Low Income" includes households with incomes up to 200% of the federal poverty level; "Mid/High Income" includes households with incomes at 200% or more of the federal poverty level.

Key Informant Input: Access to Healthcare Services

The following chart outlines key informants' perceptions of the severity of *Access to Healthcare Services* as a problem in the community:

Perceptions of Access to Healthcare Services as a Problem in the Community (Key Informants, 2018)



Sources: • PRC Online Key Informant Survey, Professional Research Consultants, Inc.
 Notes: • Asked of all respondents.

Top Concerns

Among those rating this issue as a "major problem," reasons related to the following:

Affordable Care/Services

Increasing cost of health care and shortage of doctors and other health professionals. – Community Leader (LaSalle Parish)

Lack of Providers

Number of physicians at the community clinics is not enough for the population. – Community Leader (LaSalle Parish)

Primary Care Services

About Primary Care

Improving health care services depends in part on ensuring that people have a usual and ongoing source of care. People with a usual source of care have better health outcomes and fewer disparities and costs. Having a primary care provider (PCP) as the usual source of care is especially important. PCPs can develop meaningful and sustained relationships with patients and provide integrated services while practicing in the context of family and community. Having a usual PCP is associated with:

- Greater patient trust in the provider
- Good patient-provider communication
- Increased likelihood that patients will receive appropriate care


Improving health care services includes increasing access to and use of evidence-based preventive services. Clinical preventive services are services that: **prevent** illness by detecting early warning signs or symptoms before they develop into a disease (primary prevention); or **detect** a disease at an earlier, and often more treatable, stage (secondary prevention).

- Healthy People 2020 (www.healthypeople.gov)

Access to Primary Care

This indicator is relevant because a shortage of health professionals contributes to access and health status issues.

- The proportion of primary care physicians in LaSalle Parish is much lower than found across the state and nation.



- Sources:
- US Department of Health & Human Services, Health Resources and Services Administration, Area Health Resource File.
 - Retrieved August 2018 from Community Commons at <http://www.chna.org>.
- Notes:
- This indicator is relevant because a shortage of health professionals contributes to access and health status issues.
 - These figures represent all primary care physicians practicing patient care, including hospital residents.

Specific Source of Ongoing Care

Having a specific source of ongoing care includes having a doctor’s office, clinic, urgent care center, walk-in clinic, health center facility, hospital outpatient clinic, HMO or prepaid group, military/VA clinic, or some other kind of place to go if one is sick or needs advice about his or her health. This resource is crucial to the concept of “patient-centered medical homes” (PCMH).

“Is there a particular place that you usually go to if you are sick or need advice about your health?”


“What kind of place is it: a medical clinic, an urgent care center/walk-in clinic, a doctor's office, a hospital emergency room, military or other VA healthcare, or some other place?”

The following chart illustrates the proportion of LaSalle Parish population with a specific source of ongoing medical care. Note that a hospital emergency room is not considered a specific source of ongoing care in this instance.

- The parish proportion is statistically comparable to the national proportion.
- There are no significant differences within the demographic groups.

Have a Specific Source of Ongoing Medical Care (LaSalle Parish, 2018)

Healthy People 2020 Target = 95.0% or Higher



Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 170]
 • US Department of Health and Human Services. Healthy People 2020. December 2010. <http://www.healthypeople.gov> [Objective AHS-5.1]


Notes: • Asked of all respondents.
 • Income categories reflect respondent's household income as a ratio to the federal poverty level (FPL) for their household size. "Low Income" includes households with incomes up to 200% of the federal poverty level; "Mid/High Income" includes households with incomes at 200% or more of the federal poverty level.

Utilization of Primary Care Services

“A routine checkup is a general physical exam, not an exam for a specific injury, illness or condition.

“About how long has it been since you last visited a doctor for a routine checkup?”

- The rate of routine exams in LaSalle Parish is comparable to regional, state, and US rates and is statistically similar to the 2002 baseline (though fluctuating over time).



Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 18]
 • Behavioral Risk Factor Surveillance System Survey Data. Atlanta, Georgia. United States Department of Health and Human Services, Centers for Disease Control and Prevention (CDC): 2016 LA data.
 • 2017 PRC National Health Survey, Professional Research Consultants, Inc.

Notes: • Asked of all respondents.


Emergency Room Utilization

“In the past 12 months, how many times have you gone to a hospital emergency room about your own health? This includes ER visits that resulted in a hospital admission.” (Responses here reflect the percentage with two or more visits in the past year.)

“What is the main reason you used the emergency room instead of going to a doctor's office or clinic?”

- Emergency room utilization in LaSalle Parish is similar to service area and national figures. The parish rate is comparable to the 2002 rate.

Have Used a Hospital Emergency Room More Than Once in the Past Year



Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Items 22-23]

• 2017 PRC National Health Survey, Professional Research Consultants, Inc.

Notes: • Asked of all respondents.

Oral Health

About Oral Health

Oral health is essential to overall health. Good oral health improves a person's ability to speak, smile, smell, taste, touch, chew, swallow, and make facial expressions to show feelings and emotions. However, oral diseases, from cavities to oral cancer, cause pain and disability for many Americans. Good self-care, such as brushing with fluoride toothpaste, daily flossing, and professional treatment, is key to good oral health. Health behaviors that can lead to poor oral health include: **tobacco use**; **excessive alcohol use**; and **poor dietary choices**.

The significant improvement in the oral health of Americans over the past 50 years is a public health success story. Most of the gains are a result of effective prevention and treatment efforts. One major success is community water fluoridation, which now benefits about 7 out of 10 Americans who get water through public water systems. However, some Americans do not have access to preventive programs. People who have the least access to preventive services and dental treatment have greater rates of oral diseases. A person's ability to access oral healthcare is associated with factors such as education level, income, race, and ethnicity.

Barriers that can limit a person's use of preventive interventions and treatments include: limited access to and availability of dental services; lack of awareness of the need for care; cost; and fear of dental procedures.

There are also social determinants that affect oral health. In general, people with lower levels of education and income, and people from specific racial/ethnic groups, have higher rates of disease. People with disabilities and other health conditions, like diabetes, are more likely to have poor oral health.

Potential strategies to address these issues include:


- Implementing and evaluating activities that have an impact on health behavior.
- Promoting interventions to reduce tooth decay, such as dental sealants and fluoride use.
- Evaluating and improving methods of monitoring oral diseases and conditions.
- Increasing the capacity of State dental health programs to provide preventive oral health services.
- Increasing the number of community health centers with an oral health component.
- Healthy People 2020 (www.healthypeople.gov)

Dental Care

“About how long has it been since you last visited a dentist or a dental clinic for any reason?”

- The rate of dental visits in LaSalle Parish is similar to the national figure.
- Seniors and lower-income adults are much less likely to have had a recent dental visit.

Have Visited a Dentist or Dental Clinic Within the Past Year (LaSalle Parish, 2018) Healthy People 2020 Target = 49.0% or Higher




Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 20]
 • US Department of Health and Human Services. Healthy People 2020. December 2010. <http://www.healthypeople.gov> [Objective OH-7]
 Notes: • Asked of all respondents.
 • Income categories reflect respondent's household income as a ratio to the federal poverty level (FPL) for their household size. "Low Income" includes households with incomes up to 200% of the federal poverty level; "Mid/High Income" includes households with incomes at 200% or more of the federal poverty level.

Key Informant Input: Oral Health

The following chart outlines key informants' perceptions of the severity of *Oral Health* as a problem in the community:

Perceptions of Oral Health as a Problem in the Community (Key Informants, 2018)



Sources: • PRC Online Key Informant Survey, Professional Research Consultants, Inc.
 Notes: • Asked of all respondents.

Top Concerns

Among those rating this issue as a "major problem," reasons related to the following:

Access to Care/Services


- Access to appropriate number of providers. Two dentists in LaSalle Parish. – Community Leader (LaSalle Parish)
- Only two practicing dentists in the area. – Social Services Provider (LaSalle Parish)

Vision Care

“When was the last time you had an eye exam in which the pupils were dilated? This would have made you temporarily sensitive to bright light.” (Responses in the following chart represent those with an eye exam within the past 2 years.)

- The proportion of parish adults receiving vision care is nearly identical to the Rapides Foundation Service area and statistically similar to the US. The parish rate has improved since 2002, though fluctuating dramatically in the interim.

Had an Eye Exam in the Past Two Years During Which the Pupils Were Dilated




Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 19]
 • 2017 PRC National Health Survey, Professional Research Consultants, Inc.
 Notes: • Asked of all respondents.

Local Resources

Perceptions of Local Healthcare Services

“How would you rate the overall health care services available to you? Would you say: excellent, very good, good, fair, or poor?”

- “Fair” and “poor” responses from LaSalle Parish are comparable to service area and national responses.




- Sources:
- 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 6]
 - 2017 PRC National Health Survey, Professional Research Consultants, Inc.
- Notes:
- Asked of all respondents.

Quality of Life

Life in Central Louisiana

“Now I would like to ask you some questions about this area in general. Would you say that the overall quality of life in Central Louisiana is: excellent, very good, good, fair, or poor?”


Rating of the Quality of Life in Central Louisiana
(LaSalle Parish, 2018)



Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 333]
Notes: • Asked of all respondents.

- Those in LaSalle Parish are much less likely than those across the service area to rate quality of life in Central Louisiana as “fair” or “poor.”

Quality of Life in Central Louisiana is “Fair” or “Poor”




Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 333]
Notes: • Asked of all respondents.

Charitable Contribution

“How often do you work as a volunteer for charitable organizations or community groups? Would you say: frequently, sometimes, seldom, or never?”


Frequency of Volunteering for Charitable Organizations or Community Groups
(LaSalle Parish, 2018)



Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 339]
Notes: • Asked of all respondents.

“How often do you contribute money to charitable organizations or community groups? Would you say: frequently, sometimes, seldom, or never?”

Frequency of Contributing Money to Charitable Organizations or Community Groups
(LaSalle Parish, 2018)




Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 341]
Notes: • Asked of all respondents.

“In the past 12 months, have you received assistance from a local program, church, or charitable organization to help meet some of your basic needs such as food, clothing, transportation, or child care? Please do not include any government-sponsored program or service in your response.”

- The proportion of those receiving assistance in the parish is similar to the RFSA proportion.

Received Assistance from a Local Program, Church, or Charitable Organization in the Past Month




Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 342]
 Notes: • Asked of all respondents.
 • In this case, assistance does not include government-sponsored programs or services.

Civic Participation

“For the last five times you were eligible to vote in a local, state, or national election, about how many times did you actually go and vote?”

- Adults in LaSalle Parish are much more likely to vote regularly than those across the service area.

Voted in Each of the Past Five Voting Opportunities [Including Local, State, and National Elections]



Sources: • 2018 PRC Community Health Survey, Professional Research Consultants, Inc. [Item 344]
 Notes: • Asked of all respondents.